

DHS Polyclonal Antibody

Catalog # AP69533

Product Information

Application	WB
Primary Accession	P49366
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40971

Additional Information

Gene ID	1725
Other Names	DHPS; DS; Deoxyhypusine synthase; DHS
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	DHPS
Synonyms	DS
Function	Catalyzes the NAD-dependent oxidative cleavage of spermidine and the subsequent transfer of the butylamine moiety of spermidine to the epsilon-amino group of a critical lysine residue of the eIF-5A precursor protein to form the intermediate deoxyhypusine residue (PubMed: 30661771). This is the first step of the post-translational modification of that lysine into an unusual amino acid residue named hypusine. Hypusination is unique to mature eIF-5A factor and is essential for its function.

Background


Catalyzes the NAD-dependent oxidative cleavage of spermidine and the subsequent transfer of the butylamine moiety of spermidine to the epsilon-amino group of a critical lysine residue of the eIF-5A precursor protein to form the intermediate deoxyhypusine residue. This is the first step of the post-translational modification of that lysine into an unusual amino acid residue named hypusine. Hypusination

is unique to mature eIF- 5A factor and is essential for its function.

Images



Western Blot analysis of various cells using DHS Polyclonal Antibody



Western Blot analysis of K562 cells using DHS Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.