

# **Dlx-5 Polyclonal Antibody**

Catalog # AP69543

### **Product Information**

**Application** WB, IHC-P, IF **Primary Accession** P56178

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW31540

#### **Additional Information**

**Gene ID** 1749

Other Names DLX5; Homeobox protein DLX-5

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other

applications. IHC-P~~N/A IF~~1:50~200

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

#### **Protein Information**

Name DLX5

**Function** Transcriptional factor involved in bone development. Acts as an immediate

early BMP-responsive transcriptional activator essential for osteoblast differentiation. Stimulates ALPL promoter activity in a RUNX2-independent manner during osteoblast differentiation. Stimulates SP7 promoter activity during osteoblast differentiation. Promotes cell proliferation by up-regulating

MYC promoter activity. Involved as a positive regulator of both

chondrogenesis and chondrocyte hypertrophy in the endochondral skeleton. Binds to the homeodomain-response element of the ALPL and SP7 promoter. Binds to the MYC promoter. Requires the 5'-TAATTA-3' consensus sequence

for DNA-binding.


**Cellular Location** Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00108}.

# **Background**

Transcriptional factor involved in bone development. Acts as an immediate early BMP-responsive

transcriptional activator essential for osteoblast differentiation. Stimulates ALPL promoter activity in a RUNX2-independent manner during osteoblast differentiation. Stimulates SP7 promoter activity during osteoblast differentiation. Promotes cell proliferation by up-regulating MYC promoter activity. Involved as a positive regulator of both chondrogenesis and chondrocyte hypertrophy in the endochondral skeleton. Binds to the homeodomain-response element of the ALPL and SP7 promoter. Binds to the MYC promoter. Requires the 5'-TAATTA-3' consensus sequence for DNA-binding.

## **Images**



Western Blot analysis of various cells using Dlx-5 Polyclonal Antibody diluted at 1:500

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.