

DNA pol λ Polyclonal Antibody

Catalog # AP69561

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q9UGP5
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63482

Additional Information

Gene ID	27343
Other Names	POLL; DNA polymerase lambda; Pol Lambda; DNA polymerase beta-2; Pol beta2; DNA polymerase kappa
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	POLL (HGNC:9184)
Function	DNA polymerase that functions in several pathways of DNA repair (PubMed: 11457865 , PubMed: 19806195 , PubMed: 20693240 , PubMed: 30250067). Involved in base excision repair (BER) responsible for repair of lesions that give rise to abasic (AP) sites in DNA (PubMed: 11457865 , PubMed: 19806195). Also contributes to DNA double- strand break repair by non-homologous end joining and homologous recombination (PubMed: 19806195 , PubMed: 20693240 , PubMed: 30250067). Has both template-dependent and template-independent (terminal transferase) DNA polymerase activities (PubMed: 10887191 , PubMed: 10982892 , PubMed: 12809503 , PubMed: 14627824 , PubMed: 15537631 , PubMed: 19806195). Also has a 5'-deoxyribose-5-phosphate lyase (dRP lyase) activity (PubMed: 11457865 , PubMed: 19806195).
Cellular Location	Nucleus. Chromosome. Note=Accumulates at sites of DNA damage


Tissue Location

Expressed in a number of tissues. Abundant in testis.


Background

DNA polymerase that functions in several pathways of DNA repair (PubMed: [11457865](#), PubMed:[19806195](#), PubMed:[20693240](#)). Involved in base excision repair (BER) responsible for repair of lesions that give rise to abasic (AP) sites in DNA (PubMed:[11457865](#), PubMed:[19806195](#)). Also contributes to DNA double-strand break repair by non-homologous end joining and homologous recombination (PubMed:[19806195](#), PubMed:[20693240](#)). Has both template-dependent and template-independent (terminal transferase) DNA polymerase activities (PubMed:[10982892](#), PubMed:[10887191](#), PubMed:[12809503](#), PubMed:[14627824](#), PubMed:[15537631](#), PubMed:[19806195](#)). Has also a 5'-deoxyribose-5- phosphate lyase (dRP lyase) activity (PubMed:[11457865](#), PubMed:[19806195](#)).

Images



Western Blot analysis of various cells using DNA pol λ Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100 (4°, overnight). High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.