10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

DREG Polyclonal Antibody

Catalog # AP69596

Product Information

Application	WB, IF, ICC, E
Primary Accession	Q86SQ4
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	136695

Additional Information

Gene ID	57211
Other Names	GPR126; DREG; VIGR; G-protein coupled receptor 126; Developmentally regulated G-protein-coupled receptor; Vascular inducible G protein-coupled receptor
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	ADGRG6 (HGNC:13841)
Function	Adhesion G protein-coupled receptor (aGPCR) for steroid hormones, such as progesterone and 17alpha-hydroxyprogesterone (17OHP) (PubMed:35394864, PubMed:39884271). Involved in many biological processes, such as myelination, sprouting angiogenesis, placenta, ear and cartilage development (By similarity). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide- binding proteins (G proteins) and modulates the activity of downstream effectors, such as adenylate cyclase (PubMed:24227709, PubMed:35394864). ADGRG6 is coupled to G(i) G alpha proteins and mediates inhibition of adenylate cyclase (PubMed:24227709, PubMed:35394864). Also able to couple to G(q) G proteins (PubMed:24227709). Involved in myelination of the peripheral nervous system: required for differentiation of promyelinating Schwann cells and for normal myelination of axons (PubMed:24227709). Also acts as a regulator of body length and bone mass (PubMed:18391950). Acts as a regulator of blood-brain barrier formation in the central nervous system vie its association

	with LRP1 and ITGB1 (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein. Note=Detected on the cell surface of activated but not resting umbilical vein.
Tissue Location	Expressed in placenta and to a lower extent in pancreas and liver. Detected in aortic endothelial cells but not in skin microvascular endothelial cells.

Background

G-protein coupled receptor which is activated by type IV collagen, a major constituent of the basement membrane (By similarity). Couples to G(i)-proteins as well as G(s)-proteins (PubMed:24227709). Essential for normal differentiation of promyelinating Schwann cells and for normal myelination of axons (PubMed:24227709). Regulates neural, cardiac and ear development via G-protein- and/or N-terminus-dependent signaling (By similarity). May act as a receptor for PRNP which may promote myelin homeostasis (By similarity).

Images



Western Blot analysis of various cells using DREG Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.