10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Dvl-2 Polyclonal Antibody

Catalog # AP69607

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	O14641
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78948

Additional Information

Gene ID	1856
Other Names	DVL2; Segment polarity protein dishevelled homolog DVL-2; Dishevelled-2; DSH homolog 2
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃


Protein Information

Name	DVL2
Function	Plays a role in the signal transduction pathways mediated by multiple Wnt genes (PubMed:24616100). Participates both in canonical and non-canonical Wnt signaling by binding to the cytoplasmic C- terminus of frizzled family members and transducing the Wnt signal to down-stream effectors. Promotes internalization and degradation of frizzled proteins upon Wnt signaling.
Cellular Location	Cell membrane {ECO:0000250|UniProtKB:Q60838}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q60838}; Cytoplasmic side {ECO:0000250|UniProtKB:Q60838}. Cytoplasm, cytosol. Cytoplasmic vesicle {ECO:0000250|UniProtKB:Q60838}. Nucleus Note=Localizes at the cell membrane upon interaction with frizzled family members and promotes their internalization. Localizes to cytoplasmic puncta (By similarity). Interaction with FOXK1 and FOXK2 induces nuclear translocation (PubMed:25805136) {ECO:0000250|UniProtKB:Q60838, ECO:0000269|PubMed:24616100, ECO:0000269|PubMed:25805136}

Background

Plays a role in the signal transduction pathways mediated by multiple Wnt genes. Participates both in canonical and non-canonical Wnt signaling by binding to the cytoplasmic C- terminus of frizzled family members and transducing the Wnt signal to down-stream effectors. Promotes internalization and degradation of frizzled proteins upon Wnt signaling.

Images



Western Blot analysis of K562 cells using Dvl-2 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.