

# Dynactin 2 Polyclonal Antibody

Catalog # AP69609

## Product Information

Application	WB, IHC-P
Primary Accession	<a href="#">Q13561</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44231

## Additional Information

Gene ID	10540
Other Names	DCTN2; DCTN50; Dynactin subunit 2; 50 kDa dynein-associated polypeptide; Dynactin complex 50 kDa subunit; DCTN-50; p50 dynamitin
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information


Name	DCTN2 ( <a href="#">HGNC:2712</a> )
Synonyms	DCTN50
Function	Part of the dynactin complex that activates the molecular motor dynein for ultra-processive transport along microtubules. In the dynactin soulder domain, binds the ACTR1A filament and acts as a molecular ruler to determine the length (By similarity). Modulates cytoplasmic dynein binding to an organelle, and plays a role in prometaphase chromosome alignment and spindle organization during mitosis. Involved in anchoring microtubules to centrosomes. May play a role in synapse formation during brain development (By similarity).
Cellular Location	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:A0A5G2QD80}

## Background

Modulates cytoplasmic dynein binding to an organelle, and plays a role in prometaphase chromosome alignment and spindle organization during mitosis. Involved in anchoring microtubules to centrosomes. May play a role in synapse formation during brain development.

## Images

---



Western Blot analysis of various cells using Dynactin 2 Polyclonal Antibody diluted at 1 : 500

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.