

EAAT1 Polyclonal Antibody

Catalog # AP69631

Product Information

Application	WB
Primary Accession	P43003
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59572

Additional Information

Gene ID	6507
Other Names	SLC1A3; EAAT1; GLAST; GLAST1; Excitatory amino acid transporter 1; Sodium-dependent glutamate/aspartate transporter 1; GLAST-1; Solute carrier family 1 member 3
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	SLC1A3 (HGNC:10941)
Function	Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed: 20477940 , PubMed: 26690923 , PubMed: 28032905 , PubMed: 28424515 , PubMed: 7521911 , PubMed: 8123008). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion (PubMed: 20477940). Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (PubMed: 20477940). Plays a redundant role in the rapid removal of released glutamate from the synaptic cleft, which is essential for terminating the postsynaptic action of glutamate (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein
Tissue Location	Detected in brain (PubMed: 7521911 , PubMed: 8123008 , PubMed: 8218410). Detected at very much lower levels in heart, lung, placenta and skeletal

muscle (PubMed:7521911, PubMed:8123008). Highly expressed in cerebellum, but also found in frontal cortex, hippocampus and basal ganglia (PubMed:7521911).

Background

Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed:[7521911](#), PubMed:[8123008](#), PubMed:[20477940](#), PubMed:[26690923](#), PubMed:[28032905](#), PubMed:[28424515](#)). Functions as a symporter that transports one amino acid molecule together with two or three Na⁽⁺⁾ ions and one proton, in parallel with the counter-transport of one K⁽⁺⁾ ion (PubMed:[20477940](#)). Mediates Cl⁽⁻⁾ flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na⁽⁺⁾ symport (PubMed:[20477940](#)). Plays a redundant role in the rapid removal of released glutamate from the synaptic cleft, which is essential for terminating the postsynaptic action of glutamate (By similarity).

Images



Western Blot analysis of various cells using EAAT1 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.