

EAAT3 Polyclonal Antibody

Catalog # AP69632

Product Information

Application	WB
Primary Accession	P43005
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57100

Additional Information

Gene ID	6505
Other Names	SLC1A1; EAAC1; EAAT3; Excitatory amino acid transporter 3; Excitatory amino-acid carrier 1; Neuronal and epithelial glutamate transporter; Sodium-dependent glutamate/aspartate transporter 3; Solute carrier family 1 member 1
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	SLC1A1 (HGNC:10939)
Function	Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed: 21123949 , PubMed: 26690923 , PubMed: 33658209 , PubMed: 7521911 , PubMed: 7914198 , PubMed: 8857541). Can also transport L-cysteine (PubMed: 21123949). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion (PubMed: 26690923 , PubMed: 33658209 , PubMed: 7521911 , PubMed: 8857541). Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (PubMed: 26690923 , PubMed: 8857541). Plays an important role in L- glutamate and L-aspartate reabsorption in renal tubuli (PubMed: 21123949). Plays a redundant role in the rapid removal of released glutamate from the synaptic cleft, which is essential for terminating the postsynaptic action of glutamate (By similarity). Contributes to glutathione biosynthesis and protection against oxidative

stress via its role in L-glutamate and L-cysteine transport (By similarity).
Negatively regulated by ARL6IP5 (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P43003}. Apical cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P43003}. Synapse, synaptosome {ECO:0000250|UniProtKB:P51906}. Early endosome membrane {ECO:0000250|UniProtKB:P51906}. Late endosome membrane {ECO:0000250|UniProtKB:P51906}. Recycling endosome membrane {ECO:0000250|UniProtKB:P51906}


Tissue Location

Expressed in all tissues tested including liver, muscle, testis, ovary, retinoblastoma cell line, neurons and brain (in which there was dense expression in substantia nigra, red nucleus, hippocampus and in cerebral cortical layers)


Background

Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed:[7914198](#), PubMed:[7521911](#), PubMed:[8857541](#), PubMed:[26690923](#), PubMed:[21123949](#)). Can also transport L-cysteine (PubMed:[21123949](#)). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion (PubMed:[7521911](#), PubMed:[8857541](#), PubMed:[26690923](#)). Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (PubMed:[8857541](#), PubMed:[26690923](#)). Plays an important role in L-glutamate and L-aspartate reabsorption in renal tubuli (PubMed:[21123949](#)). Plays a redundant role in the rapid removal of released glutamate from the synaptic cleft, which is essential for terminating the postsynaptic action of glutamate (By similarity). Negatively regulated by ARL6IP5 (By similarity).

Images



Western Blot analysis of various cells using EAAT3 Polyclonal Antibody



Western Blot analysis of 293 cells using EAAT3 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.