

# EDG-5 Polyclonal Antibody

Catalog # AP69650

## Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	<a href="#">O95136</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38867

## Additional Information

Gene ID	9294
Other Names	S1PR2; EDG5; Sphingosine 1-phosphate receptor 2; S1P receptor 2; S1P2; Endothelial differentiation G-protein coupled receptor 5; Sphingosine 1-phosphate receptor Edg-5; S1P receptor Edg-5
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

Name	S1PR2
Synonyms	EDG5
Function	Receptor for the lysosphingolipid sphingosine 1-phosphate (S1P) (PubMed: <a href="#">10617617</a> , PubMed: <a href="#">25274307</a> ). S1P is a bioactive lysophospholipid that elicits diverse physiological effects on most types of cells and tissues (PubMed: <a href="#">10617617</a> ). When expressed in rat HTC4 hepatoma cells, is capable of mediating S1P-induced cell proliferation and suppression of apoptosis (PubMed: <a href="#">10617617</a> ). Receptor for the chemokine-like protein FAM19A5 (PubMed: <a href="#">29453251</a> ). Mediates the inhibitory effect of FAM19A5 on vascular smooth muscle cell proliferation and migration (By similarity). In lymphoid follicles, couples the binding of S1P to the activation of GNA13 and downstream inhibition of AKT activation leading to suppression of germinal center (GC) B cell growth and migration outside the GC niche.
Cellular Location	Cell membrane; Multi-pass membrane protein


## Background

---

Receptor for the lysosphingolipid sphingosine 1- phosphate (S1P) (PubMed: [10617617](#)). S1P is a bioactive lysophospholipid that elicits diverse physiological effects on most types of cells and tissues (PubMed:[10617617](#)). When expressed in rat HTC4 hepatoma cells, is capable of mediating S1P-induced cell proliferation and suppression of apoptosis (PubMed:[10617617](#)). Receptor for the chemokine-like protein FAM19A5 (PubMed:[29453251](#)). Mediates the inhibitory effect of FAM19A5 on vascular smooth muscle cell proliferation and migration (By similarity).

## Images

---



Western Blot analysis of various cells using EDG-5 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.