

EDG-6 Polyclonal Antibody

Catalog # AP69651

Product Information

Application	WB, IF, ICC, E
Primary Accession	O95977
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41623

Additional Information

Gene ID	8698
Other Names	S1PR4; EDG6; Sphingosine 1-phosphate receptor 4; S1P receptor 4; S1P4; Endothelial differentiation G-protein coupled receptor 6; Sphingosine 1-phosphate receptor Edg-6; S1P receptor Edg-6
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	S1PR4
Synonyms	EDG6
Function	G protein-coupled receptor highly expressed in immune cells, where it regulates immune response and cytokine production. Functions as a receptor for the lysosphingolipid sphingosine-1-phosphate (S1P). Upon S1P binding, promotes regulatory T-cell differentiation and enhances fatty acid oxidation, through activation of the NRF2/PPARA signaling pathway (By similarity). Modulates also M1 macrophage activation through interaction with FPR2 and the JNK signaling, contributing to the inflammatory response (By similarity). In addition, facilitates early neutrophil mobilization and vascular activation during inflammation, promoting lymphocyte recruitment to draining lymph nodes and supporting the development of germinal centers for an effective adaptive immune response (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein.

Tissue Location

Specifically expressed in fetal and adult lymphoid and hematopoietic tissue as well as in lung. Considerable level of expression in adult and fetal spleen as well as adult peripheral leukocytes and lung. Lower expression in adult thymus, lymph node, bone marrow, and appendix as well as in fetal liver, thymus, and lung

Background

Receptor for the lysosphingolipid sphingosine 1- phosphate (S1P). S1P is a bioactive lysophospholipid that elicits diverse physiological effect on most types of cells and tissues. May be involved in cell migration processes that are specific for lymphocytes.

Images



Western Blot analysis of various cells using EDG-6 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.