

EFP Polyclonal Antibody

Catalog # AP69661

Product Information

Application	WB, IHC-P
Primary Accession	Q14258
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70973

Additional Information

Gene ID	7706
Other Names	TRIM25; EFP; RNF147; ZNF147; E3 ubiquitin/ISG15 ligase TRIM25; Estrogen-responsive finger protein; RING finger protein 147; Tripartite motif-containing protein 25; Ubiquitin/ISG15-conjugating enzyme TRIM25; Zinc finger protein 147
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	TRIM25
Synonyms	EFP {ECO:0000303 PubMed:8248217}, RNF147
Function	Functions as a ubiquitin E3 ligase and as an ISG15 E3 ligase (PubMed: 16352599). Involved in innate immune defense against viruses by mediating ubiquitination of RIGI and IFIH1 (PubMed: 17392790 , PubMed: 29357390 , PubMed: 30193849 , PubMed: 31710640 , PubMed: 33849980 , PubMed: 36045682). Mediates 'Lys-63'-linked polyubiquitination of the RIGI N-terminal CARD-like region and may play a role in signal transduction that leads to the production of interferons in response to viral infection (PubMed: 17392790 , PubMed: 23950712). Mediates 'Lys-63'-linked polyubiquitination of IFIH1 (PubMed: 30193849). Promotes ISGylation of 14-3-3 sigma (SFN), an adapter protein implicated in the regulation of a large spectrum signaling pathway (PubMed: 16352599 , PubMed: 17069755). Mediates estrogen action in various target organs (PubMed: 22452784). Mediates the ubiquitination and subsequent

proteasomal degradation of ZFHX3 (PubMed:[22452784](#)). Plays a role in promoting the restart of stalled replication forks via interaction with the KHDC3L-OOEP scaffold and subsequent ubiquitination of BLM, resulting in the recruitment and retainment of BLM at DNA replication forks (By similarity). Plays an essential role in the antiviral activity of ZAP/ZC3HAV1; an antiviral protein which inhibits the replication of certain viruses. Mechanistically, mediates 'Lys-63'-linked polyubiquitination of ZAP/ZC3HAV1 that is required for its optimal binding to target mRNA (PubMed:[28060952](#), PubMed:[28202764](#)). Also mediates the ubiquitination of various substrates implicated in stress granule formation, nonsense-mediated mRNA decay, nucleoside synthesis and mRNA translation and stability (PubMed:[36067236](#)).

Cellular Location

Cytoplasm. Cytoplasm, Stress granule. Nucleus
{ECO:0000250|UniProtKB:Q61510}


Tissue Location

Expressed in breast tumors (at protein level). Ubiquitous.

Background

Functions as a ubiquitin E3 ligase and as an ISG15 E3 ligase (PubMed: [16352599](#)). Involved in innate immune defense against viruses by mediating ubiquitination of DDX58 and IFIH1 (PubMed:[17392790](#), PubMed:[30193849](#)). Mediates 'Lys-63'-linked polyubiquitination of the DDX58 N-terminal CARD-like region which is crucial for triggering the cytosolic signal transduction that leads to the production of interferons in response to viral infection (PubMed:[17392790](#)). Mediates 'Lys-63'-linked polyubiquitination of IFIH1 (PubMed:[30193849](#)). Promotes ISGylation of 14-3-3 sigma (SFN), an adapter protein implicated in the regulation of a large spectrum signaling pathway (PubMed:[16352599](#), PubMed:[17069755](#)). Mediates estrogen action in various target organs (PubMed:[22452784](#)). Mediates the ubiquitination and subsequent proteasomal degradation of ZFHX3 (PubMed:[22452784](#)).

Images



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.