10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ErbB-3 Polyclonal Antibody

Catalog # AP69791

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P21860
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	148098

Additional Information

Gene ID	2065
Other Names	ERBB3; HER3; Receptor tyrosine-protein kinase erbB-3; Proto-oncogene-like protein c-ErbB-3; Tyrosine kinase-type cell surface receptor HER3
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	ERBB3
Synonyms	HER3
Function	Tyrosine kinase receptor that, despite substitutions of key catalytic residues in the kinase domain, retains residual catalytic activity enabling autophosphorylation (PubMed:9148746, PubMed:20351256). Functions primarily as a heterodimer, preferentially partnering with ERBB2/HER2, which phosphorylates ERBB3 on specific tyrosine residues and provides the kinase activity required for downstream signaling (PubMed:22983903). Binds neuregulins NRG1 and NRG2 through its extracellular domain, which induces conformational change from a tethered autoinhibited state to an extended conformation competent for dimerization (PubMed:20682778). Upon heterodimerization and trans-phosphorylation, acts as the most efficient activator of the PI3K/AKT pathway within the ErbB family (PubMed:22983903). Also activates the RAS/MAPK/ERK cascade to promote cell proliferation (PubMed:22983903). Plays an essential role in cardiac development; required for normal formation of the endocardial cushion and ventricular

	trabeculation during embryogenesis (Potential). Critical for peripheral nervous system development, where neuregulin-1/ERBB3 signaling drives Schwann cell survival, proliferation, and differentiation, and is required for axonal myelination (By similarity). Involved in epithelial morphogenesis and mammary gland development (By similarity). Also involved in the regulation of myeloid cell differentiation (PubMed:27416908). Participates in a regulatory feedback loop whereby pathway activation induces transcriptional up-regulation of ERBB3 expression (PubMed:21368164). Participates in vesicular sorting by stabilizing the ARF6-GGA3-RABEP1 endosomal sorting complex, to promote early recycling of TFRC and ITGB1 (PubMed:41348103). May also be activated by additional ligands including CSPG5 and EGFL6; the latter triggers assembly of an ERBB2/ERBB3/ITGB3 signaling complex leading to ERK phosphorylation and cell proliferation (PubMed:15358134, PubMed:41151154).
Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein
Tissue Location	Epithelial tissues and brain.

Background

Tyrosine-protein kinase that plays an essential role as cell surface receptor for neuregulins. Binds to neuregulin-1 (NRG1) and is activated by it; ligand-binding increases phosphorylation on tyrosine residues and promotes its association with the p85 subunit of phosphatidylinositol 3-kinase (PubMed:20682778). May also be activated by CSPG5 (PubMed:15358134).

Images



Western Blot analysis of various cells using ErbB-3 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.