

# FANCD2 Polyclonal Antibody

Catalog # AP69858

## Product Information

Application	WB, IHC-P
Primary Accession	<a href="#">Q9BXW9</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	164128

## Additional Information

Gene ID	2177
Other Names	FANCD2; FACD; Fanconi anemia group D2 protein; Protein FACD2
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

Name	FANCD2
Synonyms	FACD
Function	Required for maintenance of chromosomal stability (PubMed: <a href="#">11239453</a> , PubMed: <a href="#">14517836</a> ). Promotes accurate and efficient pairing of homologs during meiosis (PubMed: <a href="#">14517836</a> ). Involved in the repair of DNA double-strand breaks, both by homologous recombination and single-strand annealing (PubMed: <a href="#">15671039</a> , PubMed: <a href="#">15650050</a> , PubMed: <a href="#">30335751</a> , PubMed: <a href="#">36385258</a> ). The FANCI-FANCD2 complex binds and scans double-stranded DNA (dsDNA) for DNA damage; this complex stalls at DNA junctions between double-stranded DNA and single-stranded DNA (By similarity). May participate in S phase and G2 phase checkpoint activation upon DNA damage (PubMed: <a href="#">15377654</a> ). Plays a role in preventing breakage and loss of missegregating chromatin at the end of cell division, particularly after replication stress (PubMed: <a href="#">15454491</a> , PubMed: <a href="#">15661754</a> ). Required for the targeting, or stabilization, of BLM to non-centromeric abnormal structures induced by replicative stress (PubMed: <a href="#">15661754</a> , PubMed: <a href="#">19465921</a> ).

Promotes BRCA2/FANCD1 loading onto damaged chromatin (PubMed:[11239454](#), PubMed:[12239151](#), PubMed:[12086603](#), PubMed:[15115758](#), PubMed:[15199141](#), PubMed:[15671039](#), PubMed:[18212739](#)). May also be involved in B-cell immunoglobulin isotype switching.

### Cellular Location

Nucleus Note=Concentrates in nuclear foci during S phase and upon genotoxic stress. At the onset of mitosis, excluded from chromosomes and diffuses into the cytoplasm, returning to the nucleus at the end of cell division. Observed in a few spots localized in pairs on the sister chromatids of mitotic chromosome arms and not centromeres, one on each chromatids. These foci coincide with common fragile sites and could be sites of replication fork stalling. The foci are frequently interlinked through BLM-associated ultra-fine DNA bridges. Following aphidicolin treatment, targets chromatid gaps and breaks


### Tissue Location

Highly expressed in germinal center cells of the spleen, tonsil, and reactive lymph nodes, and in the proliferating basal layer of squamous epithelium of tonsil, esophagus, oropharynx, larynx and cervix. Expressed in cytotrophoblastic cells of the placenta and exocrine cells of the pancreas (at protein level). Highly expressed in testis, where expression is restricted to maturing spermatocytes


## Background

Required for maintenance of chromosomal stability. Promotes accurate and efficient pairing of homologs during meiosis. Involved in the repair of DNA double-strand breaks, both by homologous recombination and single-strand annealing. May participate in S phase and G2 phase checkpoint activation upon DNA damage. Plays a role in preventing breakage and loss of missegregating chromatin at the end of cell division, particularly after replication stress. Required for the targeting, or stabilization, of BLM to non-centromeric abnormal structures induced by replicative stress. Promotes BRCA2/FANCD1 loading onto damaged chromatin. May also be involved in B-cell immunoglobulin isotype switching.

## Images



Western Blot analysis of various cells using FANCD2 Polyclonal Antibody diluted at 1 : 500 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).



Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100(4°, overnight). High-pressure and temperature Tris-EDTA, pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.