

FOP Polyclonal Antibody

Catalog # AP69919

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	O95684
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43065

Additional Information

Gene ID	11116
Other Names	FGFR1OP; FOP; FGFR1 oncogene partner
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	CEP43 (HGNC:17012)
Synonyms	FGFR1OP, FOP
Function	Required for anchoring microtubules to the centrosomes (PubMed: 16314388 , PubMed: 28659385). Required for ciliation (PubMed: 28625565 , PubMed: 28659385).
Cellular Location	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm, cytoskeleton, cilium basal body. Note=Associated with gamma-tubulin (PubMed:16314388). Localizes on both mother and daughter centrioles (PubMed:28428259, PubMed:28625565). Localizes to an axial position on the mother centriole (PubMed:28625565). Localizes to the distal end of the centriole partly on the subdistal appendage region (PubMed:28659385).
Tissue Location	Ubiquitous. Highly expressed in heart, liver, muscle, kidney, intestine, colon, adrenal gland, prostate, testis, and pancreas.

Background

Required for anchoring microtubules to the centrosomes (PubMed: [16314388](#), PubMed:[28659385](#)).
Required for ciliation (PubMed:[28625565](#), PubMed:[28659385](#)).

Images



Western Blot analysis of various cells using FOP
Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.