

# GABAB R1 Polyclonal Antibody

Catalog # AP70011

## Product Information

---

Application	WB, IHC-P
Primary Accession	<a href="#">Q9UBS5</a>
Reactivity	Human, Mouse, Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	108320

## Additional Information

---

Gene ID	2550
Other Names	GABBR1; GPRC3A; Gamma-aminobutyric acid type B receptor subunit 1; GABA-B receptor 1; GABA-B-R1; GABA-BR1; GABABR1; Gb1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	GABBR1
Synonyms	GPRC3A
Function	Component of a heterodimeric G-protein coupled receptor for GABA, formed by GABBR1 and GABBR2 (PubMed: <a href="#">15617512</a> , PubMed: <a href="#">18165688</a> , PubMed: <a href="#">22660477</a> , PubMed: <a href="#">24305054</a> , PubMed: <a href="#">36103875</a> , PubMed: <a href="#">9872316</a> , PubMed: <a href="#">9872744</a> ). Within the heterodimeric GABA receptor, only GABBR1 seems to bind agonists, while GABBR2 mediates coupling to G proteins (PubMed: <a href="#">18165688</a> ). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase (PubMed: <a href="#">10075644</a> , PubMed: <a href="#">10773016</a> , PubMed: <a href="#">10906333</a> , PubMed: <a href="#">24305054</a> , PubMed: <a href="#">9872744</a> ). Signaling inhibits adenylate cyclase, stimulates phospholipase A2, activates potassium channels, inactivates voltage-dependent calcium-channels and modulates inositol phospholipid hydrolysis (PubMed: <a href="#">10075644</a> ). Calcium is required for high affinity binding to GABA (By similarity). Plays a critical role in the fine- tuning of inhibitory synaptic transmission (PubMed: <a href="#">9844003</a> ). Pre- synaptic GABA receptor

inhibits neurotransmitter release by down-regulating high-voltage activated calcium channels, whereas postsynaptic GABA receptor decreases neuronal excitability by activating a prominent inwardly rectifying potassium (Kir) conductance that underlies the late inhibitory postsynaptic potentials (PubMed:[10075644](#), PubMed:[22660477](#), PubMed:[9844003](#), PubMed:[9872316](#), PubMed:[9872744](#)). Not only implicated in synaptic inhibition but also in hippocampal long-term potentiation, slow wave sleep, muscle relaxation and antinociception (Probable). Activated by (-)-baclofen, cgp27492 and blocked by phaclofen (PubMed:[24305054](#), PubMed:[9844003](#), PubMed:[9872316](#)).

### Cellular Location

Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane {ECO:0000250|UniProtKB:Q9Z0U4}; Multi-pass membrane protein. Cell projection, dendrite {ECO:0000250|UniProtKB:Q9Z0U4}. Note=Colocalizes with ATF4 in hippocampal neuron dendritic membranes (By similarity). Coexpression of GABBR1 and GABBR2 is required for GABBR1 maturation and transport to the plasma membrane (PubMed:15617512). {ECO:0000250|UniProtKB:Q9Z0U4, ECO:0000269|PubMed:15617512}

### Tissue Location

Highly expressed in brain (PubMed:9753614, PubMed:9844003, PubMed:9872744). Weakly expressed in heart, small intestine and uterus. Isoform 1A: Mainly expressed in granular cell and molecular layer (PubMed:9844003). Isoform 1B: Mainly expressed in Purkinje cells (PubMed:9844003). Isoform 1E: Predominantly expressed in peripheral tissues as kidney, lung, trachea, colon, small intestine, stomach, bone marrow, thymus and mammary gland (PubMed:10906333)

## Background


---

Component of a heterodimeric G-protein coupled receptor for GABA, formed by GABBR1 and GABBR2 (PubMed:[9872316](#), PubMed:[9872744](#), PubMed:[15617512](#), PubMed:[18165688](#), PubMed:[22660477](#), PubMed:[24305054](#)). Within the heterodimeric GABA receptor, only GABBR1 seems to bind agonists, while GABBR2 mediates coupling to G proteins (PubMed:[18165688](#)). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase (PubMed:[10906333](#), PubMed:[10773016](#), PubMed:[10075644](#), PubMed:[9872744](#), PubMed:[24305054](#)). Signaling inhibits adenylate cyclase, stimulates phospholipase A2, activates potassium channels, inactivates voltage-dependent calcium-channels and modulates inositol phospholipid hydrolysis (PubMed:[10075644](#)). Calcium is required for high affinity binding to GABA (By similarity). Plays a critical role in the fine-tuning of inhibitory synaptic transmission (PubMed:[9844003](#)). Pre-synaptic GABA receptor inhibits neurotransmitter release by down-regulating high-voltage activated calcium channels, whereas postsynaptic GABA receptor decreases neuronal excitability by activating a prominent inwardly rectifying potassium (Kir) conductance that underlies the late inhibitory postsynaptic potentials (PubMed:[9844003](#), PubMed:[9872316](#), PubMed:[10075644](#), PubMed:[9872744](#), PubMed:[22660477](#)). Not only implicated in synaptic inhibition but also in hippocampal long-term potentiation, slow wave sleep, muscle relaxation and antinociception (Probable). Activated by (-)-baclofen, cgp27492 and blocked by phaclofen (PubMed:[9844003](#), PubMed:[9872316](#), PubMed:[24305054](#)).

## Images

---

Western Blot analysis of various cells using GABAB R1  
Polyclonal Antibody diluted at 1 : 500



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.