

Gasdermin C Polyclonal Antibody

Catalog # AP70038

Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | Q9BYG8 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 57692 |

Additional Information

| | |
|--------------------|--|
| Gene ID | 56169 |
| Other Names | GSDMC; MLZE; Gasdermin-C; Melanoma-derived leucine zipper-containing extranuclear factor |
| Dilution | WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. |
| Format | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide. |
| Storage Conditions | -20°C |


Protein Information

| | |
|-------------------|--|
| Name | GSDMC {ECO:0000303 PubMed:17350798, ECO:0000312 HGNC:HGNC:7151} |
| Function | [Gasdermin-C]: This form constitutes the precursor of the pore-forming protein: upon cleavage, the released N-terminal moiety (Gasdermin-C, N-terminal) binds to membranes and forms pores, triggering pyroptosis. |
| Cellular Location | [Gasdermin-C]: Cytoplasm, cytosol |
| Tissue Location | Expressed mainly in trachea and spleen (PubMed:11223543). In the esophagus, expressed in differentiating cells and probably in differentiated cells. Also detected in gastric epithelium (PubMed:19051310). |

Background

The N-terminal moiety promotes pyroptosis. May be acting by homooligomerizing within the membrane and forming pores (PubMed:[27281216](#)). The physiological relevance of this observation is unknown (Probable).

Images



Western Blot analysis of various cells using Gasdermin C
Polyclonal Antibody diluted at 1 : 500

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.