

GATA-5 Polyclonal Antibody

Catalog # AP70047

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q9BWX5
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41299

Additional Information

Gene ID	140628
Other Names	GATA5; Transcription factor GATA-5; GATA-binding factor 5
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	GATA5 (HGNC:15802)
Function	Transcription factor required during cardiovascular development (PubMed: 23289003). Plays an important role in the transcriptional program(s) that underlies smooth muscle cell diversity (By similarity). Binds to the functionally important CEF-1 nuclear protein binding site in the cardiac-specific slow/cardiac troponin C transcriptional enhancer (PubMed: 25543888).
Cellular Location	Nucleus.

Background

Transcription factor required during cardiovascular development (PubMed: [23289003](#)). Plays an important role in the transcriptional program(s) that underlies smooth muscle cell diversity (By similarity). Binds to the functionally important CEF-1 nuclear protein binding site in the cardiac-specific slow/cardiac troponin C transcriptional enhancer (PubMed:[25543888](#)).

Images



Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.