

MMP-9 Polyclonal Antibody

Catalog # AP70075

Product Information

Application WB, IHC-P, IF
Primary Accession P14780
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 78458

Additional Information

Gene ID 4318

Other Names MMP9; CLG4B; Matrix metalloproteinase-9; MMP-9; 92 kDa gelatinase; 92 kDa

type IV collagenase; Gelatinase B; GELB

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other

applications. IHC-P~~N/A IF~~1:50~200

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name MMP9

Synonyms CLG4B

Function Matrix metalloproteinase that plays an essential role in local proteolysis of

the extracellular matrix and in leukocyte migration (PubMed: 12879005, PubMed: 1480034, PubMed: 2551898). Could play a role in bone osteoclastic

resorption (By similarity). Cleaves KiSS1 at a Gly-|-Leu bond

(PubMed:<u>12879005</u>). Cleaves NINJ1 to generate the Secreted ninjurin-1 form (PubMed:<u>32883094</u>). Cleaves type IV and type V collagen into large C-terminal three quarter fragments and shorter N- terminal one quarter fragments (PubMed:<u>1480034</u>). Degrades fibronectin but not laminin or Pz-peptide.

Cellular Location Secreted, extracellular space, extracellular matrix


Tissue Location Detected in neutrophils (at protein level) (PubMed:7683678). Produced by

normal alveolar macrophages and granulocytes.


Background

May play an essential role in local proteolysis of the extracellular matrix and in leukocyte migration. Could play a role in bone osteoclastic resorption. Cleaves KiSS1 at a Gly-|-Leu bond. Cleaves type IV and type V collagen into large C-terminal three quarter fragments and shorter N-terminal one quarter fragments. Degrades fibronectin but not laminin or Pz-peptide.

Images



Western Blot analysis of various cells using MMP-9 Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of HepG2 cells using MMP-9 Polyclonal Antibody diluted at 1:1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.