

GK1 Polyclonal Antibody

Catalog # AP70093

Product Information

Application	WB, IF
Primary Accession	P32189
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61245

Additional Information

Gene ID	2710
Other Names	GK; Glycerol kinase; GK; Glycerokinase; ATP:glycerol 3-phosphotransferase
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	GK (HGNC:4289)
Function	Kinase that plays a key role in glycerol metabolism, catalyzing its phosphorylation to produce sn-glycerol 3-phosphate. Sn- glycerol 3-phosphate is a crucial intermediate in various metabolic pathways, such as the synthesis of glycerolipids and triglycerides, glycogenesis, glycolysis and gluconeogenesis.
Cellular Location	Mitochondrion outer membrane; Single-pass membrane protein. Nucleus. Cytoplasm, cytosol. Note=Glycerol kinase activity is more cytosolic in some tissues. It probably represents the expression of isoforms lacking a transmembrane domain [Isoform 4]: Cytoplasm, cytosol. Note=In adult tissues, such as liver the glycerol kinase activity is more cytosolic. It probably represents the expression of this isoform which lacks a transmembrane domain
Tissue Location	[Isoform 2]: Widely expressed in fetal and adult tissues. [Isoform 4]: The sole isoform expressed in adult liver and kidney.

Background

Key enzyme in the regulation of glycerol uptake and metabolism.

Images



Western Blot analysis of various cells using GK1
Polyclonal Antibody diluted at 1 : 2000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.