

Hexim1 Polyclonal Antibody

Catalog # AP70312

Product Information

Application	WB, IHC-P, IF
Primary Accession	O94992
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40623

Additional Information

Gene ID	10614
Other Names	HEXIM1; CLP1; EDG1; HIS1; MAQ1; Protein HEXIM1; Cardiac lineage protein 1; Estrogen down-regulated gene 1 protein; Hexamethylene bis-acetamide-inducible protein 1; Menage a quatre protein 1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	HEXIM1
Synonyms	CLP1, EDG1, HIS1, MAQ1
Function	Transcriptional regulator which functions as a general RNA polymerase II transcription inhibitor (PubMed: 14580347 , PubMed: 15201869 , PubMed: 15713661). Core component of the 7SK RNP complex: in cooperation with 7SK snRNA sequesters P-TEFb in a large inactive 7SK snRNP complex preventing RNA polymerase II phosphorylation and subsequent transcriptional elongation (PubMed: 12832472 , PubMed: 14580347 , PubMed: 15201869 , PubMed: 15713661). May also regulate NF-kappa-B, ESR1, NR3C1 and CIITA-dependent transcriptional activity (PubMed: 15940264 , PubMed: 15941832 , PubMed: 17088550). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed: 28712728).

Cellular Location	Nucleus. Cytoplasm. Note=Binds alpha-importin and is mostly nuclear (PubMed:16362050)
Tissue Location	Ubiquitously expressed with higher expression in placenta. HEXIM1 and HEXIM2 are differentially expressed. Expressed in endocrine tissues.

Background

Transcriptional regulator which functions as a general RNA polymerase II transcription inhibitor (PubMed:[14580347](#), PubMed:[15713661](#), PubMed:[15201869](#)). In cooperation with 7SK snRNA sequesters P-TEFb in a large inactive 7SK snRNP complex preventing RNA polymerase II phosphorylation and subsequent transcriptional elongation (PubMed:[12832472](#), PubMed:[14580347](#), PubMed:[15713661](#), PubMed:[15201869](#)). May also regulate NF-kappa-B, ESR1, NR3C1 and CIITA-dependent transcriptional activity (PubMed:[15940264](#), PubMed:[15941832](#), PubMed:[17088550](#)). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:[28712728](#)).

Images



Western Blot analysis of RAT-HEART cells using Hexim1 Polyclonal Antibody diluted at 1 : 2000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.