

HGK Polyclonal Antibody

Catalog # AP70313

Product Information

Application WB, IF, ICC, E **Primary Accession** 095819

Reactivity Human, Mouse

HostRabbitClonalityPolyclonalCalculated MW142101

Additional Information

Gene ID 9448

Other Names MAP4K4; HGK; KIAA0687; NIK; Mitogen-activated protein kinase kinase kinase

kinase 4; HPK/GCK-like kinase HGK; MAPK/ERK kinase kinase kinase 4; MEK

kinase kinase 4; MEKKK 4; Nck-interacting kinase

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000.

ELISA: 1/40000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A

E~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name MAP4K4 (HGNC:6866)

Synonyms HGK, KIAA0687, NIK

Function Serine/threonine kinase that plays a role in the response to environmental

stress and cytokines such as TNF-alpha. Appears to act upstream of the JUN N-terminal pathway (PubMed: 9890973). Activator of the Hippo signaling pathway which plays a pivotal role in organ size control and tumor

suppression by restricting proliferation and promoting apoptosis. MAP4Ks act in parallel to and are partially redundant with STK3/MST2 and STK4/MST2 in the phosphorylation and activation of LATS1/2, and establish MAP4Ks as components of the expanded Hippo pathway (PubMed: 26437443).

Phosphorylates SMAD1 on Thr- 322 (PubMed: 21690388).


Cellular Location Cytoplasm.

Tissue Location Widely expressed. Isoform 5 is abundant in the brain. Isoform 4 is

Background

Serine/threonine kinase that may play a role in the response to environmental stress and cytokines such as TNF-alpha. Appears to act upstream of the JUN N-terminal pathway. Phosphorylates SMAD1 on Thr-322.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.