

HGK Polyclonal Antibody

Catalog # AP70313

Product Information

Application	WB, IF
Primary Accession	O95819
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	142101

Additional Information

Gene ID	9448
Other Names	MAP4K4; HGK; KIAA0687; NIK; Mitogen-activated protein kinase kinase kinase 4; HPK/GCK-like kinase HGK; MAPK/ERK kinase kinase 4; MEK kinase kinase 4; MEKKK 4; Nck-interacting kinase
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	MAP4K4 (HGNC:6866)
Synonyms	HGK, KIAA0687, NIK
Function	Serine/threonine kinase that plays a role in the response to environmental stress and cytokines such as TNF-alpha. Appears to act upstream of the JUN N-terminal pathway (PubMed: 9890973). Activator of the Hippo signaling pathway which plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. MAP4Ks act in parallel to and are partially redundant with STK3/MST2 and STK4/MST2 in the phosphorylation and activation of LATS1/2, and establish MAP4Ks as components of the expanded Hippo pathway (PubMed: 26437443). Phosphorylates SMAD1 on Thr- 322 (PubMed: 21690388).
Cellular Location	Cytoplasm.
Tissue Location	Widely expressed. Isoform 5 is abundant in the brain. Isoform 4 is predominant in the liver, skeletal muscle and placenta.

Background

Serine/threonine kinase that may play a role in the response to environmental stress and cytokines such as TNF-alpha. Appears to act upstream of the JUN N-terminal pathway. Phosphorylates SMAD1 on Thr-322.

Images



Western Blot analysis of various cells using HGK Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.