

Histone deacetylase 10 Polyclonal Antibody

Catalog # AP70329

Product Information

| | |
|-------------------|---------------------------|
| Application | WB, IHC-P |
| Primary Accession | Q969S8 |
| Reactivity | Human, Mouse, Rat, Monkey |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 71445 |

Additional Information

| | |
|--------------------|---|
| Gene ID | 83933 |
| Other Names | HDAC10; Histone deacetylase 10; HD10 |
| Dilution | WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A |
| Format | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide. |
| Storage Conditions | -20°C |


Protein Information

| | |
|-------------------|---|
| Name | HDAC10 |
| Function | <p>Polyamine deacetylase (PDAC), which acts preferentially on N(8)-acetylspermidine, and also on acetylcadaverine and acetylputrescine (PubMed:28516954). Exhibits attenuated catalytic activity toward N(1),N(8)-diacetylspermidine and very low activity, if any, toward N(1)-acetylspermidine (PubMed:28516954). Histone deacetylase activity has been observed in vitro (PubMed:11677242, PubMed:11726666, PubMed:11739383, PubMed:11861901). Has also been shown to be involved in MSH2 deacetylation (PubMed:26221039). The physiological relevance of protein/histone deacetylase activity is unclear and could be very weak (PubMed:28516954). May play a role in the promotion of late stages of autophagy, possibly autophagosome- lysosome fusion and/or lysosomal exocytosis in neuroblastoma cells (PubMed:23801752, PubMed:29968769). May play a role in homologous recombination (PubMed:21247901). May promote DNA mismatch repair (PubMed:26221039).</p> |
| Cellular Location | Cytoplasm. Nucleus Note=Excluded from nucleoli. |
| Tissue Location | Widely expressed with high levels in liver and kidney. |

Background

Polyamine deacetylase (PDAC), which acts preferentially on N(8)-acetylspermidine, and also on acetylcadaverine and acetylputrescine (PubMed:[28516954](#)). Exhibits attenuated catalytic activity toward N(1),N(8)-diacetylspermidine and very low activity, if any, toward N(1)-acetylspermidine (PubMed:[28516954](#)). Histone deacetylase activity has been observed in vitro (PubMed:[11861901](#), PubMed:[11726666](#), PubMed:[11677242](#), PubMed:[11739383](#)). Has also been shown to be involved in MSH2 deacetylation (PubMed:[26221039](#)). The physiological relevance of protein/histone deacetylase activity is unclear and could be very weak (PubMed:[28516954](#)). May play a role in the promotion of late stages of autophagy, possibly autophagosome-lysosome fusion and/or lysosomal exocytosis in neuroblastoma cells (PubMed:[23801752](#), PubMed:[29968769](#)). May play a role in homologous recombination (PubMed:[21247901](#)). May promote DNA mismatch repair (PubMed:[26221039](#)).

Images



Western Blot analysis of various cells using Histone deacetylase 10 Polyclonal Antibody diluted at 1 : 1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.