

IKKy Polyclonal Antibody

Catalog # AP70488

Product Information

Application	WB, IHC-P, IF
Primary Accession	Q9Y6K9
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48198

Additional Information

Gene ID	8517
Other Names	IKBKG; FIP3; NEMO; NF-kappa-B essential modulator; NEMO; FIP-3; IκB kinase-associated protein 1; IKKAP1; Inhibitor of nuclear factor kappa-B kinase subunit gamma; I-kappa-B kinase subunit gamma; IKK-gamma; IKKG; IκB kinase subunit gamma; NF
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	IKBKG (HGNC:5961)
Synonyms	FIP3, NEMO
Function	Regulatory subunit of the IKK core complex which phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor (PubMed: 14695475 , PubMed: 20724660 , PubMed: 21518757 , PubMed: 9751060). Its binding to scaffolding polyubiquitin plays a key role in IKK activation by multiple signaling receptor pathways (PubMed: 16547522 , PubMed: 18287044 , PubMed: 19033441 , PubMed: 19185524 , PubMed: 21606507 , PubMed: 27777308 , PubMed: 33567255). Can recognize and bind both 'Lys-63'-linked and linear polyubiquitin upon cell stimulation, with a much higher affinity for linear polyubiquitin (PubMed: 16547522 , PubMed: 18287044 , PubMed: 19033441 , PubMed: 19185524 , PubMed: 21606507 , PubMed: 27777308). Could be implicated in NF-kappa-B-mediated protection


from cytokine toxicity. Essential for viral activation of IRF3 (PubMed:[19854139](#)). Involved in TLR3- and IFIH1-mediated antiviral innate response; this function requires 'Lys- 27'-linked polyubiquitination (PubMed:[20724660](#)).

Cellular Location	Cytoplasm. Nucleus Note=Sumoylated NEMO accumulates in the nucleus in response to genotoxic stress.
Tissue Location	Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

Background

Regulatory subunit of the IKK core complex which phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor. Its binding to scaffolding polyubiquitin seems to play a role in IKK activation by multiple signaling receptor pathways. However, the specific type of polyubiquitin recognized upon cell stimulation (either 'Lys-63'-linked or linear polyubiquitin) and its functional importance is reported conflictingly. Also considered to be a mediator for TAX activation of NF-kappa-B. Could be implicated in NF-kappa-B- mediated protection from cytokine toxicity. Essential for viral activation of IRF3. Involved in TLR3- and IFIH1-mediated antiviral innate response; this function requires 'Lys-27'-linked polyubiquitination.

Images



Western Blot analysis of various cells using IKKγ Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.