

IL-15R α Polyclonal Antibody

Catalog # AP70498

Product Information

Application	WB
Primary Accession	Q13261
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28233

Additional Information

Gene ID	3601
Other Names	IL15RA; Interleukin-15 receptor subunit alpha; IL-15 receptor subunit alpha; IL-15R-alpha; IL-15RA; CD antigen CD215
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	IL15RA
Function	High-affinity receptor for interleukin-15 (PubMed: 8530383). Can signal both in cis and trans where IL15R from one subset of cells presents IL15 to neighboring IL2RG-expressing cells (By similarity). In neutrophils, binds and activates kinase SYK in response to IL15 stimulation (PubMed: 15123770). In neutrophils, required for IL15- induced phagocytosis in a SYK-dependent manner (PubMed: 15123770). Expression of different isoforms may alter or interfere with signal transduction (PubMed: 10480910).
Cellular Location	Membrane; Single- pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein. Cell surface. Note=Mainly found associated with the nuclear membrane [Isoform 6]; Endoplasmic reticulum membrane; Single-pass type I membrane protein. Golgi apparatus membrane; Single- pass type I membrane protein. Cytoplasmic vesicle membrane; Single-pass type I membrane protein. Membrane; Single-pass type I membrane protein Note=Isoform 5, isoform 6, isoform 7 and isoform 8 are associated with endoplasmic reticulum, Golgi and cytoplasmic vesicles, but not with the nuclear membrane [Isoform 8]; Endoplasmic reticulum

membrane; Single-pass type I membrane protein. Golgi apparatus membrane; Single-pass type I membrane protein. Cytoplasmic vesicle membrane; Single-pass type I membrane protein. Membrane; Single-pass type I membrane protein Note=Isoform 5, isoform 6, isoform 7 and isoform 8 are associated with endoplasmic reticulum, Golgi and cytoplasmic vesicles, but not with the nuclear membrane


Tissue Location

Expressed in neutrophils (at protein level) (PubMed:15123770). Expressed in fetal brain with higher expression in the hippocampus and cerebellum than in cortex and thalamus (PubMed:12114302). Higher levels of soluble sIL-15RA form in comparison with membrane-bound forms is present in all brain structures (PubMed:12114302). Isoforms 1, 3, 4, 5, 6, 7, 8 and 9: Widely expressed (PubMed:10480910, PubMed:8530383).


Background

High-affinity receptor for interleukin-15 (PubMed: [8530383](#)). Can signal both in cis and trans where IL15R from one subset of cells presents IL15 to neighboring IL2RG- expressing cells (By similarity). In neutrophils, binds and activates kinase SYK in response to IL15 stimulation (PubMed:[15123770](#)). In neutrophils, required for IL15-induced phagocytosis in a SYK-dependent manner (PubMed:[15123770](#)). Expression of different isoforms may alter or interfere with signal transduction (PubMed:[10480910](#)).

Images



Western Blot analysis of COLO cells using IL-15Rα Polyclonal Antibody



Western blot analysis of various lysis using IL-15Rα Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.