10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Intestinal Cell Kinase Polyclonal Antibody

Catalog # AP70558

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q9UPZ9
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	71427

Additional Information

Gene ID	22858
Other Names	ICK; KIAA0936; Serine/threonine-protein kinase ICK; Intestinal cell kinase; hICK; Laryngeal cancer kinase 2; LCK2; MAK-related kinase; MRK
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	CILK1
Synonyms	ICK, KIAA0936
Function	Required for ciliogenesis (PubMed: 24797473). Phosphorylates KIF3A (By similarity). Involved in the control of ciliary length (PubMed:24853502). Regulates the ciliary localization of SHH pathway components as well as the localization of IFT components at ciliary tips (By similarity). May play a key role in the development of multiple organ systems and particularly in cardiac development (By similarity). Regulates intraflagellar transport (IFT) speed and negatively regulates cilium length in a cAMP and mTORC1 signaling-dependent manner and this regulation requires its kinase activity (By similarity).
Cellular Location	Nucleus. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q62726}. Cell projection, cilium. Cytoplasm, cytoskeleton, cilium basal body {ECO:0000250|UniProtKB:Q9JKV2}. Note=Also found at the ciliary tip (PubMed:24797473). Nuclear localization has been observed with a GFP-

	tagged construct in transfected HeLa cells (PubMed:12103360, PubMed:19185282).
Tissue Location	Expressed in heart, brain, placenta, pancreas, thymus, prostate, testis, ovary, small intestine and colon, with highest levels in placenta and testis. Not detected in spleen. Also expressed in many cancer cell lines.

Background

Required for ciliogenesis (PubMed: 24797473). Phosphorylates KIF3A (By similarity). Involved in the control of ciliary length (PubMed:24853502). Regulates the ciliary localization of SHH pathway components as well as the localization of IFT components at ciliary tips (By similarity). May play a key role in the development of multiple organ systems and particularly in cardiac development (By similarity). Regulates intraflagellar transport (IFT) speed and negatively regulates cilium length in a cAMP and mTORC1 signaling-dependent manner and this regulation requires its kinase activity (By similarity).

Images



Western Blot analysis of 3T3 cells using Intestinal Cell Kinase Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.