

IRAK-M Polyclonal Antibody

Catalog # AP70576

Product Information

Application	WB, IHC-P, IF
Primary Accession	Q9Y616
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67767

Additional Information

Gene ID	11213
Other Names	IRAK3; Interleukin-1 receptor-associated kinase 3; IRAK-3; IL-1 receptor-associated kinase M; IRAK-M
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	IRAK3 {ECO:0000312 EMBL:AAH57800.1}
Function	Putative inactive protein kinase which regulates signaling downstream of immune receptors including IL1R and Toll-like receptors (PubMed: 10383454 , PubMed: 29686383). Inhibits dissociation of IRAK1 and IRAK4 from the Toll-like receptor signaling complex by either inhibiting the phosphorylation of IRAK1 and IRAK4 or stabilizing the receptor complex (By similarity). Upon IL33-induced lung inflammation, positively regulates expression of IL6, CSF3, CXCL2 and CCL5 mRNAs in dendritic cells (PubMed: 29686383).
Cellular Location	Cytoplasm. Nucleus. Note=In dendritic cells, translocates into the nucleus upon IL33 stimulation. {ECO:0000250 UniProtKB:Q8K4B2}
Tissue Location	Expressed in eosinophils, dendritic cells and/or monocytes (at protein level) (PubMed:29686383). Expressed predominantly in peripheral blood lymphocytes (PubMed:10383454)

Background

Inhibits dissociation of IRAK1 and IRAK4 from the Toll- like receptor signaling complex by either inhibiting the phosphorylation of IRAK1 and IRAK4 or stabilizing the receptor complex.

Images



Western Blot analysis of LOVO cells using IRAK-M Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.