

# Ku-86 Polyclonal Antibody

Catalog # AP70687

## Product Information

---

Application	WB, IHC-P
Primary Accession	<a href="#">P13010</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	82705

## Additional Information

---

Gene ID	7520
Other Names	XRCC5; G22P2; X-ray repair cross-complementing protein 5; 86 kDa subunit of Ku antigen; ATP-dependent DNA helicase 2 subunit 2; ATP-dependent DNA helicase II 80 kDa subunit; CTC box-binding factor 85 kDa subunit; CTC85; CTCBF; DNA repair pr
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	XRCC5
Synonyms	G22P2
Function	Single-stranded DNA-dependent ATP-dependent helicase that plays a key role in DNA non-homologous end joining (NHEJ) by recruiting DNA-PK to DNA (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> ). Required for double-strand break repair and V(D)J recombination (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> ). Also has a role in chromosome translocation (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> ). The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> ). It works in the 3'-5' direction (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> ).

During NHEJ, the XRCC5-XRCC6 dimer performs the recognition step: it recognizes and binds to the broken ends of the DNA and protects them from further resection (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[7957065](#), PubMed:[8621488](#)). Binding to DNA may be mediated by XRCC6 (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[7957065](#), PubMed:[8621488](#)). The XRCC5-XRCC6 dimer acts as a regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20383123](#), PubMed:[7957065](#), PubMed:[8621488](#)). The XRCC5-XRCC6 dimer is probably involved in stabilizing broken DNA ends and bringing them together (PubMed:[12145306](#), PubMed:[20383123](#), PubMed:[7957065](#), PubMed:[8621488](#)). The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step (PubMed:[12145306](#), PubMed:[20383123](#), PubMed:[7957065](#), PubMed:[8621488](#)). The XRCC5-XRCC6 dimer probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5'-deoxyribose-5-phosphate at an abasic site near double-strand breaks (PubMed:[20383123](#)). XRCC5 probably acts as the catalytic subunit of 5'-dRP activity, and allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined (PubMed:[20383123](#)). The XRCC5-XRCC6 dimer together with APEX1 acts as a negative regulator of transcription (PubMed:[8621488](#)). In association with NAA15, the XRCC5-XRCC6 dimer binds to the osteocalcin promoter and activates osteocalcin expression (PubMed:[12145306](#)). As part of the DNA-PK complex, involved in the early steps of ribosome assembly by promoting the processing of precursor rRNA into mature 18S rRNA in the small-subunit processome (PubMed:[32103174](#)). Binding to U3 small nucleolar RNA, recruits PRKDC and XRCC5/Ku86 to the small-subunit processome (PubMed:[32103174](#)). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:[28712728](#)).

#### Cellular Location

Nucleus. Nucleus, nucleolus Chromosome


## Background

---


Single-stranded DNA-dependent ATP-dependent helicase. Has a role in chromosome translocation. The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner. It works in the 3'-5' direction. Binding to DNA may be mediated by XRCC6. Involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination. The XRCC5/6 dimer acts as regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold. The XRCC5/6 dimer is probably involved in stabilizing broken DNA ends and bringing them together (PubMed:[12145306](#), PubMed:[20383123](#), PubMed:[7957065](#), PubMed:[8621488](#)). The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step. In association with NAA15, the XRCC5/6 dimer binds to the osteocalcin promoter and activates osteocalcin expression (PubMed:[20383123](#)). The XRCC5/6 dimer probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5'-deoxyribose-5-phosphate at an abasic site near double-strand breaks. XRCC5 probably acts as the catalytic subunit of 5'-dRP activity, and allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined. The XRCC5/6 dimer together with APEX1 acts as a negative regulator of transcription (PubMed:[8621488](#)). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway.

## Images


---



Western Blot analysis of various cells using Ku-86 Polyclonal Antibody



Western Blot analysis of A549 cells using Ku-86 Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.