10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

KV3.1 Polyclonal Antibody

Catalog # AP70693

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P48547
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57942

Additional Information

Gene ID	3746
Other Names	KCNC1; Potassium voltage-gated channel subfamily C member 1; NGK2; Voltage-gated potassium channel subunit Kv3.1; Voltage-gated potassium channel subunit Kv4
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	KCNC1 {ECO:0000303|PubMed:8449507, ECO:0000312|HGNC:HGNC:6233}
Function	Voltage-gated potassium channel that opens in response to the voltage difference across the membrane and through which potassium ions pass in accordance with their electrochemical gradient (PubMed:25401298, PubMed:35840580). The mechanism is time-dependent and inactivation is slow (By similarity). Plays an important role in the rapid repolarization of fast-firing brain neurons (By similarity). Can form functional homotetrameric channels and heterotetrameric channels that contain variable proportions of KCNC2, and possibly other family members as well (By similarity). Contributes to fire sustained trains of very brief action potentials at high frequency in pallidal neurons (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein. Cell projection, axon {ECO:0000250|UniProtKB:P25122}. Presynaptic cell membrane {ECO:0000250|UniProtKB:P25122}. Note=Localizes in parallel fiber membranes, distributed on the perisynaptic and extrasynaptic membranes

away from the active zones. {ECO:0000250|UniProtKB:P25122}


Background

Voltage-gated potassium channel that plays an important role in the rapid repolarization of fast-firing brain neurons. The channel opens in response to the voltage difference across the membrane, forming a potassium-selective channel through which potassium ions pass in accordance with their electrochemical gradient (PubMed:25401298). Can form functional homotetrameric channels and heterotetrameric channels that contain variable proportions of KCNC2, and possibly other family members as well. Contributes to fire sustained trains of very brief action potentials at high frequency in pallidal neurons.

Images



Western Blot analysis of various cells using KV3.1 Polyclonal Antibody



Western Blot analysis of K562 cells using KV3.1 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.