

KV4.1 Polyclonal Antibody

Catalog # AP70696

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q9NSA2
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	71330

Additional Information

Gene ID	3750
Other Names	KCND1; Potassium voltage-gated channel subfamily D member 1; Voltage-gated potassium channel subunit Kv4.1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	KCND1 (HGNC:6237)
Function	A-type voltage-gated potassium channel that mediates transmembrane potassium transport in excitable membranes in the brain (PubMed: 15454437). Mediates A-type current I(SA) in suprachiasmatic nucleus (SCN) neurons. Exhibits a low-threshold A-type current with a hyperpolarized steady-state inactivation midpoint and the recovery process was steeply voltage-dependent, with recovery being markedly faster at more negative potentials. May regulates repetitive firing rates in the suprachiasmatic nucleus (SCN) neurons and circadian rhythms in neuronal excitability and behavior. Contributes to the regulation of the circadian rhythm of action potential firing in suprachiasmatic nucleus neurons, which regulates the circadian rhythm of locomotor activity. The regulatory subunit KCNIP1 modulates the kinetics of channel inactivation, increases the current amplitudes and accelerates recovery from inactivation, shifts activation in a depolarizing direction (By similarity). The regulatory subunit DPP10 decreases the voltage sensitivity of the inactivation channel gating (PubMed: 15454437).

Cellular Location	Cell membrane {ECO:0000250 UniProtKB:Q9NZV8}; Multi-pass membrane protein {ECO:0000250 UniProtKB:Q9NZV8}
Tissue Location	Widely expressed. Highly expressed in brain, in particular in cerebellum and thalamus; detected at lower levels in the other parts of the brain.

Background

Pore-forming (alpha) subunit of voltage-gated rapidly inactivating A-type potassium channels. May contribute to I(To) current in heart and I(Sa) current in neurons. Channel properties are modulated by interactions with other alpha subunits and with regulatory subunits.

Images



Western Blot analysis of 3T3 cells using KV4.1 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.