

# Lad Polyclonal Antibody

Catalog # AP70701

#### **Product Information**

Application WB, IHC-P
Primary Accession Q9NP31
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 42934

#### **Additional Information**

**Gene ID** 9047

Other Names SH2D2A; SCAP; TSAD; VRAP; SH2 domain-containing protein 2A; SH2

domain-containing adapter protein; T cell-specific adapter protein; TSAd;

VEGF receptor-associated protein

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

#### **Protein Information**

Name SH2D2A

**Synonyms** SCAP, TSAD, VRAP

**Function** Could be a T-cell-specific adapter protein involved in the control of T-cell

activation. May play a role in the CD4-p56-LCK- dependent signal transduction pathway. Could also play an important role in normal and pathological angiogenesis. Could be an adapter protein that facilitates and regulates interaction of KDR with effector proteins important to endothelial cell survival

and proliferation.

Cellular Location Cytoplasm.

**Tissue Location** Expression limited to tissues of the immune system and, in particular,


activated T-cells. Expressed in peripheral blood leukocytes, thymus and spleen. Much lower expression or undetectable, in brain, placenta, skeletal muscle, prostate, testis, ovary, small intestine, and colon. Expressed at low levels in unstimulated T-cells, but not expressed in normal resting or activated

B-cells. According to PubMed:10692392, expression is not restricted to activated T-cells, but strongly expressed in blood cell lineages, the endothelium and other cell and tissue types, such as heart, lung, and liver

## **Background**

Could be a T-cell-specific adapter protein involved in the control of T-cell activation. May play a role in the CD4-p56- LCK-dependent signal transduction pathway. Could also play an important role in normal and pathological angiogenesis. Could be an adapter protein that facilitates and regulates interaction of KDR with effector proteins important to endothelial cell survival and proliferation.

### **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.