10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MAGI-2 Polyclonal Antibody

Catalog # AP70810

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q86UL8
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	158754

Additional Information

Gene ID	9863
Other Names	MAGI2; ACVRINP1; AIP1; KIAA0705; Membrane-associated guanylate kinase; WW and PDZ domain-containing protein 2; Atrophin-1-interacting protein 1; AIP-1; Atrophin-1-interacting protein A; Membrane-associated guanylate kinase inverted 2; MAGI-
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	MAGI2
Synonyms	ACVRINP1, AIP1, KIAA0705
Function	Seems to act as a scaffold molecule at synaptic junctions by assembling neurotransmitter receptors and cell adhesion proteins (By similarity). Plays a role in nerve growth factor (NGF)-induced recruitment of RAPGEF2 to late endosomes and neurite outgrowth (By similarity). May play a role in regulating activin-mediated signaling in neuronal cells (By similarity). Enhances the ability of PTEN to suppress AKT1 activation (PubMed:10760291). Plays a role in receptor- mediated clathrin-dependent endocytosis which is required for ciliogenesis (By similarity).
Cellular Location	Cytoplasm. Late endosome. Synapse, synaptosome. Cell membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q9WVQ1}. Cell

	projection, cilium {ECO:0000250|UniProtKB:Q9WVQ1}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole {ECO:0000250|UniProtKB:Q9WVQ1}. Photoreceptor inner segment {ECO:0000250|UniProtKB:Q9WVQ1}. Cell projection, cilium, photoreceptor outer segment {ECO:0000250|UniProtKB:Q9WVQ1}. Note=Localized diffusely in the cytoplasm before nerve growth factor (NGF) stimulation Recruited to late endosomes after NGF stimulation. Membrane-associated in synaptosomes (By similarity).
Tissue Location	Specifically expressed in brain.

Background

Seems to act as scaffold molecule at synaptic junctions by assembling neurotransmitter receptors and cell adhesion proteins. May play a role in regulating activin-mediated signaling in neuronal cells. Enhances the ability of PTEN to suppress AKT1 activation. Plays a role in nerve growth factor (NGF)-induced recruitment of RAPGEF2 to late endosomes and neurite outgrowth.

Images



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.