10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MBD3 Polyclonal Antibody

Catalog # AP70854

Product Information

Application	WB, E, IHC-P
Primary Accession	O95983
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32844

Additional Information

Gene ID	53615
Other Names	MBD3; Methyl-CpG-binding domain protein 3; Methyl-CpG-binding protein MBD3
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. E~~N/A IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	MBD3
Function	Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (PubMed:12124384, PubMed:16428440, PubMed:28977666). Acts as transcriptional repressor and plays a role in gene silencing (PubMed:10947852, PubMed:18644863). Does not bind to methylated DNA by itself (PubMed:12124384, PubMed:16428440). Binds to a lesser degree DNA containing unmethylated CpG dinucleotides (PubMed:24307175). Recruits histone deacetylases and DNA methyltransferases.
Cellular Location	Nucleus. Chromosome. Note=Nuclear, in discrete foci. Detected on chromatin, at promoter regions of active genes

Background

Acts as transcriptional repressor and plays a role in gene silencing. Does not bind to DNA by itself (PubMed:12124384). Binds to DNA with a preference for sites containing methylated CpG dinucleotides (in

vitro). Binds to a lesser degree DNA containing unmethylated CpG dinucleotides (PubMed:24307175). Recruits histone deacetylases and DNA methyltransferases.

Images



Western Blot analysis of various cells using MBD3 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.