

MDFI Polyclonal Antibody

Catalog # AP70871

Product Information

Application	WB, E
Primary Accession	Q99750
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25029

Additional Information

Gene ID	4188
Other Names	MDFI; MyoD family inhibitor; Myogenic repressor I-mf
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	MDFI
Function	Inhibits the transactivation activity of the Myod family of myogenic factors and represses myogenesis (By similarity). Acts by associating with Myod family members and retaining them in the cytoplasm by masking their nuclear localization signals (By similarity). Can also interfere with the DNA-binding activity of Myod family members (By similarity). Plays an important role in trophoblast and chondrogenic differentiation (By similarity). Regulates the transcriptional activity of TCF7L1/TCF3 by interacting directly with TCF7L1/TCF3 and preventing it from binding DNA (By similarity). Binds to the axin complex, resulting in an increase in the level of free beta-catenin (By similarity). Affects axin regulation of the WNT and JNK signaling pathways (By similarity). Regulates the activity of mechanosensitive Piezo channel (PubMed: 37590348).
Cellular Location	Nucleus. Cytoplasm {ECO:0000250 UniProtKB:P70331}

Background

Inhibits the transactivation activity of the Myod family of myogenic factors and represses myogenesis. Acts by associating with Myod family members and retaining them in the cytoplasm by masking their nuclear localization signals. Can also interfere with the DNA-binding activity of Myod family members. Plays an important role in trophoblast and chondrogenic differentiation. Regulates the transcriptional activity of TCF7L1/TCF3 by interacting directly with TCF7L1/TCF3 and preventing it from binding DNA. Binds to the axin complex, resulting in an increase in the level of free beta-catenin. Affects axin regulation of the WNT and JNK signaling pathways (By similarity).

Images



Western Blot analysis of various cells using MDF1 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.