

# MNDA Polyclonal Antibody

Catalog # AP70986

## **Product Information**

Application	WB, IHC-P
Primary Accession	<u>P41218</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	45836

#### **Additional Information**

Gene ID	4332
Other Names	MNDA; Myeloid cell nuclear differentiation antigen
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

# **Protein Information**

Name	MNDA
Function	May act as a transcriptional activator/repressor in the myeloid lineage. Plays a role in the granulocyte/monocyte cell-specific response to interferon. Stimulates the DNA binding of the transcriptional repressor protein YY1.
Cellular Location	Nucleus. Cytoplasm. Note=Uniformly distributed throughout the interphase cell nucleus. Associates with chromatin
Tissue Location	Expressed constitutively in cells of the myeloid lineage. Found in promyelocyte stage cells as well as in all other stage cells including peripheral blood monocytes and granulocytes. Also appears in myeloblast cells in some cases of acute myeloid Leukemia

## Background

May act as a transcriptional activator/repressor in the myeloid lineage. Plays a role in the granulocyte/monocyte cell- specific response to interferon. Stimulates the DNA binding of the transcriptional repressor protein YY1.

### Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.