

Moesin/Ezrin/Radixin Polyclonal Antibody

Catalog # AP70996

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P26038 , P35241 , P15311
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67820

Additional Information


Gene ID	4478
Other Names	MSN; Moesin; Membrane-organizing extension spike protein; RDX; Radixin; EZR; VIL2; Ezrin; Cytovillin; Villin-2; p81
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	MSN (HGNC:7373)
Function	Ezrin-radixin-moesin (ERM) family protein that connects the actin cytoskeleton to the plasma membrane and thereby regulates the structure and function of specific domains of the cell cortex. Tethers actin filaments by oscillating between a resting and an activated state providing transient interactions between moesin and the actin cytoskeleton (PubMed:10212266). Once phosphorylated on its C-terminal threonine, moesin is activated leading to interaction with F-actin and cytoskeletal rearrangement (PubMed:10212266). These rearrangements regulate many cellular processes, including cell shape determination, membrane transport, and signal transduction (PubMed:12387735, PubMed:15039356). The role of moesin is particularly important in immunity acting on both T and B-cells homeostasis and self-tolerance, regulating lymphocyte egress from lymphoid organs (PubMed:9298994, PubMed:9616160). Modulates phagolysosomal biogenesis in macrophages (By similarity). Also participates in immunologic synapse formation (PubMed:27405666).

Cellular Location	Cell membrane; Peripheral membrane protein {ECO:0000250 UniProtKB:P26041}; Cytoplasmic side {ECO:0000250 UniProtKB:P26041}. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:P26041}. Apical cell membrane {ECO:0000250 UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250 UniProtKB:P26041}; Cytoplasmic side {ECO:0000250 UniProtKB:P26041}. Cell projection, microvillus membrane {ECO:0000250 UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250 UniProtKB:P26041}; Cytoplasmic side {ECO:0000250 UniProtKB:P26041}. Cell projection, microvillus {ECO:0000250 UniProtKB:P26041}. Note=Phosphorylated form is enriched in microvilli-like structures at apical membrane. Increased cell membrane localization of both phosphorylated and non-phosphorylated forms seen after thrombin treatment (By similarity). Localizes at the uropods of T lymphoblasts. {ECO:0000250 UniProtKB:P26041, ECO:0000269 PubMed:18586956, ECO:0000269 PubMed:9298994} Tissue Location In all tissues and cultured cells studied.
---------------------------------	---

Images



Western Blot analysis of 3T3 cells using Moesin/Ezrin/Radixin Polyclonal Antibody diluted at 1 : 2000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.