

Myotubularin Polyclonal Antibody

Catalog # AP71135

Product Information

Application	WB
Primary Accession	<u>Q13496</u>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	69932

Additional Information

Gene ID	4534
Other Names	MTM1; CG2; Myotubularin
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	MTM1 (<u>HGNC:7448</u>)
Synonyms	CG2
Function	Lipid phosphatase which dephosphorylates phosphatidylinositol 3-monophosphate (PI3P) and phosphatidylinositol 3,5-bisphosphate (PI(3,5)P2) (PubMed: <u>10900271</u> , PubMed: <u>11001925</u> , PubMed: <u>12646134</u> , PubMed: <u>14722070</u>). Has also been shown to dephosphorylate phosphotyrosine- and phosphoserine-containing peptides (PubMed: <u>9537414</u>). Negatively regulates EGFR degradation through regulation of EGFR trafficking from the late endosome to the lysosome (PubMed: <u>14722070</u>). Plays a role in vacuolar formation and morphology. Regulates desmin intermediate filament assembly and architecture (PubMed: <u>21135508</u>). Plays a role in mitochondrial morphology and positioning (PubMed: <u>21135508</u>). Required for skeletal muscle maintenance but not for myogenesis (PubMed: <u>21135508</u>). In skeletal muscles, stabilizes MTMR12 protein levels (PubMed: <u>23818870</u>).
Cellular Location	Cytoplasm. Cell membrane; Peripheral membrane protein. Cell projection, filopodium. Cell projection, ruffle. Late endosome. Cytoplasm, myofibril, sarcomere {ECO:0000250 UniProtKB:Q9Z2C5}. Note=Localizes as a dense

cytoplasmic network (PubMed:11001925). Also localizes to the plasma membrane, including plasma membrane extensions such as filopodia and ruffles (PubMed:12118066). Predominantly located in the cytoplasm following interaction with MTMR12 (PubMed:12847286). Recruited to the late endosome following EGF stimulation (PubMed:14722070). In skeletal muscles, co-localizes with MTMR12 in the sarcomere (By similarity) {ECO:0000250|UniProtKB:Q9Z2C5, ECO:0000269|PubMed:11001925, ECO:0000269|PubMed:12118066, ECO:0000269|PubMed:12847286, ECO:0000269|PubMed:14722070}

Background

Lipid phosphatase which dephosphorylates phosphatidylinositol 3-monophosphate (PI3P) and phosphatidylinositol 3,5-bisphosphate (PI(3,5)P2) (PubMed:<u>11001925</u>, PubMed:<u>10900271</u>, PubMed:<u>12646134</u>, PubMed:<u>14722070</u>). Has also been shown to dephosphorylate phosphotyrosine- and phosphoserine-containing peptides (PubMed:<u>9537414</u>). Negatively regulates EGFR degradation through regulation of EGFR trafficking from the late endosome to the lysosome (PubMed:<u>14722070</u>). Plays a role in vacuolar formation and morphology. Regulates desmin intermediate filament assembly and architecture (PubMed:<u>21135508</u>). Plays a role in mitochondrial morphology and positioning (PubMed:<u>21135508</u>). Required for skeletal muscle maintenance but not for myogenesis (PubMed:<u>21135508</u>). In skeletal muscles, stabilizes MTMR12 protein levels (PubMed:<u>23818870</u>).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.