10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

NAB2 Polyclonal Antibody

Catalog # AP71158

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q15742
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56594

Additional Information

Gene ID	4665
Other Names	NAB2; MADER; NGFI-A-binding protein 2; EGR-1-binding protein 2; Melanoma-associated delayed early response protein; Protein MADER
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃


Protein Information

Name	NAB2
Synonyms	MADER
Function	Acts as a transcriptional repressor for zinc finger transcription factors EGR1 and EGR2. Isoform 2 lacks repression ability (By similarity).
Cellular Location	Nucleus. Note=Isoform 2 is not localized to the nucleus.
Tissue Location	Widely expressed at low levels. Highly expressed in melanoma cell lines

Background

Acts as a transcriptional repressor for zinc finger transcription factors EGR1 and EGR2. Isoform 2 lacks repression ability (By similarity).

Images



Western Blot analysis of various cells using NAB2 Polyclonal Antibody diluted at 1：1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.