

NAB2 Polyclonal Antibody

Catalog # AP71158

Product Information

Application WB, IHC-P Primary Accession Q15742

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Calculated MW 56594

Additional Information

Gene ID 4665

Other Names NAB2; MADER; NGFI-A-binding protein 2; EGR-1-binding protein 2;

Melanoma-associated delayed early response protein; Protein MADER

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name NAB2

Synonyms MADER

Function Acts as a transcriptional repressor for zinc finger transcription factors EGR1

and EGR2. Isoform 2 lacks repression ability (By similarity).


Cellular Location Nucleus. Note=Isoform 2 is not localized to the nucleus.

Tissue Location Widely expressed at low levels. Highly expressed in melanoma cell lines

Background

Acts as a transcriptional repressor for zinc finger transcription factors EGR1 and EGR2. Isoform 2 lacks repression ability (By similarity).

Images



Western Blot analysis of various cells using NAB2 Polyclonal Antibody diluted at 1: 1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.