

# Neddylin Polyclonal Antibody

Catalog # AP71209

## Product Information

Application	WB, IHC-P
Primary Accession	<a href="#">Q15843</a>
Reactivity	Human, Mouse, Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	9072

## Additional Information

Gene ID	4738
Other Names	NEDD8; NEDD8; Neddylin; Neural precursor cell expressed developmentally down-regulated protein 8; NEDD-8; Ubiquitin-like protein Nedd8
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

Name	NEDD8 {ECO:0000303   PubMed:9694792, ECO:0000312   HGNC:HGNC:7732}
Function	Ubiquitin-like protein which plays an important role in cell cycle control and embryogenesis via its conjugation to a limited number of cellular proteins, such as cullins or p53/TP53 (PubMed: <a href="#">10318914</a> , PubMed: <a href="#">10597293</a> , PubMed: <a href="#">11953428</a> , PubMed: <a href="#">14690597</a> , PubMed: <a href="#">15242646</a> , PubMed: <a href="#">9694792</a> , PubMed: <a href="#">38605244</a> , PubMed: <a href="#">38316879</a> ). Attachment of NEDD8 to cullins is critical for the recruitment of E2 to the cullin-RING- based E3 ubiquitin-protein ligase complex, thus facilitating polyubiquitination and proteasomal degradation of cyclins and other regulatory proteins (PubMed: <a href="#">10318914</a> , PubMed: <a href="#">10597293</a> , PubMed: <a href="#">11953428</a> , PubMed: <a href="#">20688984</a> , PubMed: <a href="#">9694792</a> , PubMed: <a href="#">38605244</a> , PubMed: <a href="#">38316879</a> ). Attachment of NEDD8 to p53/TP53 inhibits p53/TP53 transcriptional activity (PubMed: <a href="#">15242646</a> ). Covalent attachment to its substrates requires prior activation by the E1 complex UBE1C-APPBP1 and linkage to the E2 enzyme UBE2M (PubMed: <a href="#">14690597</a> ).
Cellular Location	Nucleus. Note=Mainly nuclear.

**Tissue Location**

Highly expressed in heart, skeletal muscle, spleen, thymus, prostate, testis, ovary, colon and leukocytes


**Background**

---

Ubiquitin-like protein which plays an important role in cell cycle control and embryogenesis. Covalent attachment to its substrates requires prior activation by the E1 complex UBE1C- APPBP1 and linkage to the E2 enzyme UBE2M. Attachment of NEDD8 to cullins activates their associated E3 ubiquitin ligase activity, and thus promotes polyubiquitination and proteasomal degradation of cyclins and other regulatory proteins.

**Images**

---



Western Blot analysis of various cells using Neddylin Polyclonal Antibody diluted at 1 : 1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.