

# Netrin-1 Polyclonal Antibody

Catalog # AP71225

## Product Information

---

<b>Application</b>	WB, IHC-P
<b>Primary Accession</b>	<a href="#">O95631</a>
<b>Reactivity</b>	Human, Mouse, Rat, Monkey
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	67748

## Additional Information

---

<b>Gene ID</b>	9423
<b>Other Names</b>	NTN1; NTN1L; Netrin-1; Epididymis tissue protein Li 131P
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	NTN1
<b>Synonyms</b>	NTN1L
<b>Function</b>	Netrins control guidance of CNS commissural axons and peripheral motor axons. Its association with either DCC or some UNC5 receptors will lead to axon attraction or repulsion, respectively. Binding to UNC5C might cause dissociation of UNC5C from polymerized TUBB3 in microtubules and thereby lead to increased microtubule dynamics and axon repulsion (PubMed: <a href="#">28483977</a> ). Involved in dorsal root ganglion axon projection towards the spinal cord (PubMed: <a href="#">28483977</a> ). It also serves as a survival factor via its association with its receptors which prevent the initiation of apoptosis. Involved in tumorigenesis by regulating apoptosis (PubMed: <a href="#">15343335</a> ).
<b>Cellular Location</b>	Secreted. Cytoplasm. Note=Mainly secreted
<b>Tissue Location</b>	Widely expressed in normal adult tissues with highest levels in heart, small intestine, colon, liver and prostate Reduced expression in brain tumors and neuroblastomas. Expressed in epididymis (at protein level).


## Background

---


Netrins control guidance of CNS commissural axons and peripheral motor axons. Its association with either DCC or some UNC5 receptors will lead to axon attraction or repulsion, respectively. It also serve as a survival factor via its association with its receptors which prevent the initiation of apoptosis. Involved in tumorigenesis by regulating apoptosis.

## Images

---



Western Blot analysis of various cells using Netrin-1  
Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of COS7 cells using Netrin-1  
Polyclonal Antibody diluted at 1 : 1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.