

# Nibrin Polyclonal Antibody

Catalog # AP71305

## Product Information

---

Application	WB, IHC-P
Primary Accession	<a href="#">O60934</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	84959

## Additional Information

---

Gene ID	4683
Other Names	NBN; NBS; NBS1; P95; Nibrin; Cell cycle regulatory protein p95; Nijmegen breakage syndrome protein 1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	NBN ( <a href="#">HGNC:7652</a> )
Function	<p>Component of the MRN complex, which plays a central role in double-strand break (DSB) repair, DNA recombination, maintenance of telomere integrity and meiosis (PubMed:<a href="#">10888888</a>, PubMed:<a href="#">15616588</a>, PubMed:<a href="#">18411307</a>, PubMed:<a href="#">18583988</a>, PubMed:<a href="#">18678890</a>, PubMed:<a href="#">19759395</a>, PubMed:<a href="#">23115235</a>, PubMed:<a href="#">28216226</a>, PubMed:<a href="#">28867292</a>, PubMed:<a href="#">9705271</a>). The MRN complex is involved in the repair of DNA double-strand breaks (DSBs) via homologous recombination (HR), an error-free mechanism which primarily occurs during S and G2 phases (PubMed:<a href="#">19759395</a>, PubMed:<a href="#">28867292</a>, PubMed:<a href="#">9705271</a>). The complex (1) mediates the end resection of damaged DNA, which generates proper single-stranded DNA, a key initial steps in HR, and is (2) required for the recruitment of other repair factors and efficient activation of ATM and ATR upon DNA damage (PubMed:<a href="#">19759395</a>, PubMed:<a href="#">9705271</a>). The MRN complex possesses single-strand endonuclease activity and double-strand-specific 3'-5' exonuclease activity, which are provided by MRE11, to initiate end resection,</p>

which is required for single-strand invasion and recombination (PubMed:[19759395](#), PubMed:[28867292](#), PubMed:[9705271](#)). Within the MRN complex, NBN acts as a protein-protein adapter, which specifically recognizes and binds phosphorylated proteins, promoting their recruitment to DNA damage sites (PubMed:[12419185](#), PubMed:[15616588](#), PubMed:[18411307](#), PubMed:[18582474](#), PubMed:[18583988](#), PubMed:[18678890](#), PubMed:[19759395](#), PubMed:[19804756](#), PubMed:[23762398](#), PubMed:[24534091](#), PubMed:[27814491](#), PubMed:[27889449](#), PubMed:[33836577](#)). Recruits MRE11 and RAD50 components of the MRN complex to DSBs in response to DNA damage (PubMed:[12419185](#), PubMed:[18411307](#), PubMed:[18583988](#), PubMed:[18678890](#), PubMed:[24534091](#), PubMed:[26438602](#)). Promotes the recruitment of PI3/PI4-kinase family members ATM, ATR, and probably DNA-PKcs to the DNA damage sites, activating their functions (PubMed:[15064416](#), PubMed:[15616588](#), PubMed:[15790808](#), PubMed:[16622404](#), PubMed:[22464731](#), PubMed:[30952868](#), PubMed:[35076389](#)). Mediates the recruitment of phosphorylated RBBP8/CtIP to DSBs, leading to cooperation between the MRN complex and RBBP8/CtIP to initiate end resection (PubMed:[19759395](#), PubMed:[27814491](#), PubMed:[27889449](#), PubMed:[33836577](#)). RBBP8/CtIP specifically promotes the endonuclease activity of the MRN complex to clear DNA ends containing protein adducts (PubMed:[27814491](#), PubMed:[27889449](#), PubMed:[30787182](#), PubMed:[33836577](#)). The MRN complex is also required for the processing of R-loops (PubMed:[31537797](#)). NBN also functions in telomere length maintenance via its interaction with TERF2: interaction with TERF2 during G1 phase preventing recruitment of DCLRE1B/Apollo to telomeres (PubMed:[10888888](#), PubMed:[28216226](#)). NBN also promotes DNA repair choice at dysfunctional telomeres: NBN phosphorylation by CDK2 promotes non-homologous end joining repair at telomeres, while unphosphorylated NBN promotes microhomology-mediated end-joining (MMEJ) repair (PubMed:[28216226](#)). Enhances AKT1 phosphorylation possibly by association with the mTORC2 complex (PubMed:[23762398](#)).

#### Cellular Location

Nucleus. Chromosome. Nucleus, PML body. Chromosome, telomere  
 Note=Localizes to discrete nuclear foci after treatment with genotoxic agents (PubMed:[10783165](#), PubMed:[26215093](#), PubMed:[26438602](#)). Localizes to DNA double-strand breaks (DSBs); recruited to DNA damage sites via association with phosphorylated proteins, such as phosphorylated H2AX, phosphorylated MDC1 and phosphorylated RAD17 (PubMed:[12419185](#), PubMed:[18411307](#), PubMed:[18582474](#), PubMed:[18583988](#), PubMed:[18678890](#), PubMed:[19338747](#), PubMed:[23115235](#), PubMed:[24534091](#), PubMed:[26438602](#)). Acetylation of 'Lys-5' of histone H2AX (H2AXK5ac) promotes NBN/NBS1 assembly at the sites of DNA damage (PubMed:[26438602](#)).

#### Tissue Location


Ubiquitous (PubMed:[9590180](#)). Expressed at high levels in testis (PubMed:[9590180](#)).

## Background

Component of the MRE11-RAD50-NBN (MRN complex) which plays a critical role in the cellular response to DNA damage and the maintenance of chromosome integrity. The complex is involved in double-strand break (DSB) repair, DNA recombination, maintenance of telomere integrity, cell cycle checkpoint control and meiosis. The complex possesses single-strand endonuclease activity and double-strand-specific 3'-5' exonuclease activity, which are provided by MRE11. RAD50 may be required to bind DNA ends and hold them in close proximity. NBN modulate the DNA damage signal sensing by recruiting PI3/PI4-kinase family members ATM, ATR, and probably DNA-PKcs to the DNA damage sites and activating their functions. It can also recruit MRE11 and RAD50 to the proximity of DSBs by an interaction with the histone H2AX. NBN also functions in telomere length maintenance by generating the 3' overhang which serves as a primer for

telomerase dependent telomere elongation. NBN is a major player in the control of intra-S-phase checkpoint and there is some evidence that NBN is involved in G1 and G2 checkpoints. The roles of NBS1/MRN encompass DNA damage sensor, signal transducer, and effector, which enable cells to maintain DNA integrity and genomic stability. Forms a complex with RBBP8 to link DNA double-strand break sensing to resection. Enhances AKT1 phosphorylation possibly by association with the mTORC2 complex.


## Images




Western Blot analysis of various cells using Nibrin Polyclonal Antibody



Western Blot analysis of HT29 cells using Nibrin Polyclonal Antibody




Immunohistochemical analysis of paraffin-embedded Human lung. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human lung. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Immunohistochemical analysis of paraffin-embedded Human lung. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature



EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.