

# NRBF-2 Polyclonal Antibody

Catalog # AP71370

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q96F24</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	32378

## Additional Information

---

<b>Gene ID</b>	29982
<b>Other Names</b>	NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	NRBF2
<b>Synonyms</b>	COPR
<b>Function</b>	May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).
<b>Cellular Location</b>	Nucleus {ECO:0000250 UniProtKB:Q9QYK3}. Cytoplasm {ECO:0000250 UniProtKB:Q9QYK3}. Cytoplasmic vesicle. Cytoplasmic vesicle, autophagosome
<b>Tissue Location</b>	Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852).


## Background

---

May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).

## Images

---



Western Blot analysis of various cells using NRBF-2 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.