

NY-CO-1 Polyclonal Antibody

Catalog # AP71398

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	O60524
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	122954

Additional Information

Gene ID	9147
Other Names	NEMF; SDCCAG1; Nuclear export mediator factor NEMF; Antigen NY-CO-1; Serologically defined colon cancer antigen 1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	NEMF {ECO:0000303 PubMed:33048237, ECO:0000312 HGNC:HGNC:10663}
Function	Key component of the ribosome quality control complex (RQC), a ribosome-associated complex that mediates the extraction of incompletely synthesized nascent chains from stalled ribosomes as well as their ubiquitin-mediated proteasomal degradation (PubMed: 25578875 , PubMed: 32726578 , PubMed: 33406423 , PubMed: 33909987). Thereby, frees 60S subunit ribosomes from the stalled translation complex and prevents the accumulation of nascent polypeptide chains that are potentially toxic for the cell (PubMed: 25578875 , PubMed: 33406423 , PubMed: 33909987). Within the RQC complex, NEMF specifically binds stalled 60S ribosomal subunits by recognizing an exposed, nascent chain-conjugated tRNA moiety and promotes the recruitment of LTN1 to stalled 60S subunits (PubMed: 25578875). Following binding to stalled 60S ribosomal subunits, NEMF mediates CAT tailing by recruiting alanine-charged tRNA to the A- site and directing the elongation of stalled nascent chains independently of mRNA or 40S subunits, leading to non-templated C- terminal alanine extensions (CAT tails)

(PubMed:[33406423](#), PubMed:[33909987](#)). Mainly recruits alanine-charged tRNAs, but can also recruit other amino acid-charged tRNAs (PubMed:[33406423](#), PubMed:[33909987](#)). CAT tailing is required to promote ubiquitination of stalled nascent chains by different E3 ubiquitin-protein ligases (PubMed:[33909987](#)). In the canonical RQC pathway (RQC-L), CAT tailing facilitates LTN1-dependent ubiquitination by exposing lysine residues that would otherwise remain buried in the ribosomal exit tunnel (By similarity). In the alternative RQC pathway (RQC-C) CAT tailing creates an C-degron mainly composed of alanine that is recognized by the CRL2(KLHDC10) and RCHY1/PIRH2 E3 ligases, leading to ubiquitination and degradation of stalled nascent chains (PubMed:[33909987](#)). NEMF may also indirectly play a role in nuclear export (PubMed:[16103875](#)).

Cellular Location

Cytoplasm, cytosol. Nucleus


Tissue Location

Expressed in brain, heart, liver, lung, spleen, and skeletal muscle. Also expressed at lower levels in stomach and testis


Background

Component of the ribosome quality control complex (RQC), a ribosome-associated complex that mediates ubiquitination and extraction of incompletely synthesized nascent chains for proteasomal degradation. NEMF is responsible for selective recognition of stalled 60S subunits by recognizing an exposed, nascent chain-conjugated tRNA moiety. NEMF is important for the stable association of LTN1 to the complex (PubMed:[25578875](#)). May indirectly play a role in nuclear export (PubMed:[16103875](#)).

Images




Western Blot analysis of various cells using NY-CO-1 Polyclonal Antibody



Western Blot analysis of 3T3 cells using NY-CO-1 Polyclonal Antibody

Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°, overnight). High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative



ctrl (right) obtained from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.