

PAR-4 Polyclonal Antibody

Catalog # AP71771

Product Information

Application WB, IF Primary Accession Q96RIO

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 41133

Additional Information

Gene ID 9002

Other Names F2RL3; PAR4; Proteinase-activated receptor 4; PAR-4; Coagulation factor II

receptor-like 3; Thrombin receptor-like 3

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000.

ELISA: 1/10000. Not yet tested in other applications. IF~~1:50~200

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name F2RL3

Synonyms PAR4

Function Receptor for activated thrombin or trypsin coupled to G proteins that

stimulate phosphoinositide hydrolysis (PubMed: 10079109). May play a role in

platelets activation (PubMed: 10079109).

Cellular Location Cell membrane; Multi-pass membrane protein.

Tissue Location Widely expressed, with highest levels in lung, pancreas, thyroid, testis and


small intestine. Not expressed in brain, kidney, spinal cord and peripheral

blood leukocytes. Also detected in platelets

Background

Receptor for activated thrombin or trypsin coupled to G proteins that stimulate phosphoinositide hydrolysis. May play a role in platelets activation.

Images



Western Blot analysis of various cells using PAR-4 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.