

PDE10A Polyclonal Antibody

Catalog # AP71810

Product Information

Application	WB, E, IHC-P
Primary Accession	Q9Y233
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	114946

Additional Information

Gene ID	10846
Other Names	PDE10A; cAMP and cAMP-inhibited cGMP 3'; 5'-cyclic phosphodiesterase 10A
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications. E~~N/A IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	PDE10A
Function	Plays a role in signal transduction by regulating the intracellular concentration of cyclic nucleotides (PubMed: 10373451 , PubMed: 10393245 , PubMed: 16330539 , PubMed: 17389385 , PubMed: 27058447). Can hydrolyze both cAMP and cGMP, but has higher affinity for cAMP and is more efficient with cAMP as substrate (PubMed: 10373451 , PubMed: 10393245 , PubMed: 17389385 , PubMed: 27058447). May play a critical role in regulating cAMP and cGMP levels in the striatum, a region of the brain that contributes to the control of movement and cognition (PubMed: 27058447).
Cellular Location	Cytoplasm, cytosol
Tissue Location	Abundant in the putamen and caudate nucleus regions of brain and testis, moderately expressed in the thyroid gland, pituitary gland, thalamus and cerebellum.

Background

Plays a role in signal transduction by regulating the intracellular concentration of cyclic nucleotides. Can hydrolyze both cAMP and cGMP, but has higher affinity for cAMP and is more efficient with cAMP as substrate. May play a critical role in regulating cAMP and cGMP levels in the striatum, a region of the brain that contributes to the control of movement and cognition.

Images



Western Blot analysis of A549 cells using PDE10A Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.