

PEPT1 Polyclonal Antibody

Catalog # AP71845

Product Information

Application	WB
Primary Accession	P46059
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78806

Additional Information

Gene ID	6564
Other Names	SLC15A1; PEPT1; Solute carrier family 15 member 1; Intestinal H(+)/peptide cotransporter; Oligopeptide transporter; small intestine isoform; Peptide transporter 1
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	SLC15A1 (HGNC:10920)
Function	Electrogenic proton-coupled amino-acid transporter that transports oligopeptides of 2 to 4 amino acids with a preference for dipeptides. Transports neutral and monovalently charged peptides with a proton to peptide stoichiometry of 1:1 or 2:1 (By similarity) (PubMed: 15521010 , PubMed: 18367661 , PubMed: 19685173 , PubMed: 26320580 , PubMed: 7896779 , PubMed: 8914574 , PubMed: 9835627). Primarily responsible for the absorption of dietary di- and tripeptides from the small intestinal lumen (By similarity). Mediates transepithelial transport of muramyl and N-formylated bacterial dipeptides contributing to recognition of pathogenic bacteria by the mucosal immune system (PubMed: 15521010 , PubMed: 9835627).
Cellular Location	Apical cell membrane; Multi-pass membrane protein. Note=Localized to the apical membrane of enterocytes
Tissue Location	Expressed in small intestine.

Background

Proton-coupled intake of oligopeptides of 2 to 4 amino acids with a preference for dipeptides. May constitute a major route for the absorption of protein digestion end-products.

Images



Western Blot analysis of MOUSE-BRAIN MOUSE-HEART
MOUSE-LUNG MOUSE-KIDNEY cells using PEPT1
Polyclonal Antibody diluted at 1 : 1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.