

PGE synthase Polyclonal Antibody

Catalog # AP71872

Product Information

Application	WB
Primary Accession	O14684
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17102

Additional Information

Gene ID	9536
Other Names	PTGES; MGST1L1; MPGES1; PGES; PIG12; Prostaglandin E synthase; Microsomal glutathione S-transferase 1-like 1; MGST1-L1; Microsomal prostaglandin E synthase 1; MPGES-1; p53-induced gene 12 protein
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	PTGES
Function	Terminal enzyme of the cyclooxygenase (COX)-2-mediated prostaglandin E2 (PGE2) biosynthetic pathway. Catalyzes the glutathione-dependent oxidoreduction of prostaglandin endoperoxide H2 (PGH2) to prostaglandin E2 (PGE2) in response to inflammatory stimuli (PubMed: 10377395 , PubMed: 10869354 , PubMed: 12244105 , PubMed: 12460774 , PubMed: 12672824 , PubMed: 18682561). Plays a key role in inflammation response, fever and pain (By similarity). Also catalyzes the oxidoreduction of endocannabinoids into prostaglandin glycerol esters and PGG2 into 15-hydroperoxy-PGE2 (PubMed: 12244105 , PubMed: 12672824). In addition, displays low glutathione transferase and glutathione- dependent peroxidase activities, toward 1-chloro-2,4-dinitrobenzene and 5-hydroperoxyicosatetraenoic acid (5-HPETE), respectively (PubMed: 12672824).
Cellular Location	Membrane; Multi-pass membrane protein. Cytoplasm, perinuclear region. Note=Colocalizes with PTGS1/COX-1 and PTGS2/COX-2 in the perinuclear

compartment

Background

Catalyzes the oxidoreduction of prostaglandin endoperoxide H2 (PGH2) to prostaglandin E2 (PGE2).

Images



Western Blot analysis of various cells using PGE synthase Polyclonal Antibody diluted at 1 : 500

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.