

# PLC $\gamma$ 2 Polyclonal Antibody

Catalog # AP71978

## Product Information

Application	WB, IHC-P
Primary Accession	<a href="#">P16885</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	147870

## Additional Information

Gene ID	5336
Other Names	PLCG2; 1-phosphatidylinositol 4; 5-bisphosphate phosphodiesterase gamma-2; Phosphoinositide phospholipase C-gamma-2; Phospholipase C-IV; PLC-IV; Phospholipase C-gamma-2; PLC-gamma-2
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


## Protein Information

Name	PLCG2 ( <a href="#">HGNC:9066</a> )
Function	The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. It is a crucial enzyme in transmembrane signaling.
Cellular Location	Membrane raft {ECO:0000250   UniProtKB:Q8CIH5}.

## Background

The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. It is a crucial enzyme in transmembrane signaling.

## Images



Western blot analysis of kb 3T3 lysis using PLC  $\gamma$ 2 antibody. Antibody was diluted at 1:1000

## Citations

---

- [A novel homozygous mutation of phospholipase C zeta leading to defective human oocyte activation and fertilization failure](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.