10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PTH Polyclonal Antibody

Catalog # AP72080

Product Information

Application	WB, IHC-P
Primary Accession	P01270
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	12861

Additional Information

Gene ID	5741
Other Names	PTH; Parathyroid hormone; PTH; Parathormone; Parathyrin
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information

Name	PTH {ECO:0000303|PubMed:35932760, ECO:0000312|HGNC:HGNC:9606}
Function	Peptide hormone secreted by parathyroid chief cells that acts as a primary endocrine regulator of extracellular calcium and phosphate homeostasis, acting primarily on bone and kidney to maintain serum calcium levels within a narrow range (PubMed:11604398, PubMed:35932760). Acts by binding and activating the G protein-coupled receptor PTH1R, engaging the receptor to activate G(s)-mediated cAMP/PKA signaling and G(q)-mediated PLC/PKC signaling (PubMed:18375760, PubMed:35932760). In bone, acts indirectly on osteoclasts by binding PTH1R on cells of the osteoblast/osteocyte lineage, where it increases expression of TNFSF11/RANKL and decreases expression of TNFRSF11B/OPG, thereby raising the RANKL/OPG ratio and stimulating osteoclast differentiation and bone resorption (PubMed:9797718, PubMed:11454517). Stimulates also [1-14C]-2-deoxy-D- glucose (2DG) transport and glycogen synthesis in osteoblastic cells (PubMed:21076856). In distal renal tubule cells, stimulates active Ca(2+) reabsorption via the adenylyl cyclase-cAMP-protein kinase A (PKA) pathway leading to the phosphorylation and activation of the calcium selective cation channel TRPV5 (PubMed:19423690). Acting on PTH1R in proximal renal tubule cells, decreases reabsorption of filtered phosphate by reducing apical abundance


of SLC34A1, through a mechanism involving PTH1R-activated PKC-mediated phosphorylation of the scaffold NHERF1, causing dissociation of the NHERF1-SLC34A1 complex and subsequent clathrin-mediated endocytosis and lysosomal degradation of the cotransporter (By similarity). Activates also transcription of CYP27B1 (25-hydroxyvitamin D3 1-alpha-hydroxylase) through a cAMP/PKA/SIK2-SIK3/CRTC2 axis, thereby increasing renal synthesis of the active vitamin D metabolite 1,25-dihydroxyvitamin D(3) (By similarity).

Cellular Location Secreted

Background

PTH elevates calcium level by dissolving the salts in bone and preventing their renal excretion. Stimulates [1-14C]-2- deoxy-D-glucose (2DG) transport and glycogen synthesis in osteoblastic cells.

Images



Western Blot analysis of various cells using PTH Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.